



National Institute for Public Health
and the Environment
Ministry of Health, Welfare and Sport



Environmental *radioactivity* in the Netherlands

Results in 2016

This report contains an erratum
d.d. 24-09-2019 on page 118



National Institute for Public Health
and the Environment
Ministry of Health, Welfare and Sport

Environmental radioactivity in the Netherlands

Results in 2016

This report contains an erratum
d.d. 24-09-2019 on page 118

RIVM Report 2018-0160

Colophon

© RIVM 2019

Parts of this publication may be reproduced, provided acknowledgement is given to the 'National Institute for Public Health and the Environment', along with the title and year of publication.

DOI 10.21945/RIVM-2018-0160

C.P. Tanzi (editor), RIVM

Contact:

Cristina Tanzi, Centre for Environmental Safety and Security
cristina.tanzi@rivm.nl



National Institute for Public Health
and the Environment
Ministry of Health, Welfare and Sport



Rijkswaterstaat
*Ministry of Infrastructure
and Water Management*



Netherlands Food and Consumer
Product Safety Authority
Ministry of Economic Affairs



RIKILT
WAGENINGEN UR



N.V. Elektriciteits-Produktiemaatschappij Zuid-Nederland EPZ

This investigation has been performed by order and for the account of the Authority for Nuclear Safety and Radiation Protection, within the framework of Project 390010: environmental monitoring of radioactivity and radiation.

This is a publication of:
**National Institute for Public Health
and the Environment**
P.O. Box 1 | 3720 BA Bilthoven
The Netherlands
www.rivm.nl/en

Synopsis

Environmental radioactivity in the Netherlands

Results in 2016

In 2016 the Netherlands met its annual European obligation to measure radioactivity in the environment and in food. All Member States of the European Union are required to perform these measurements each year under the terms of the Euratom Treaty of 1957. The Netherlands complied with the recommendations, as established in 2000, to perform these measurements in a uniform manner.

The results on radioactivity in the environment are reported to the European Commission by the National Institute for Public Health and the Environment (RIVM) on behalf of the competent authority in the Netherlands.

The measurements represent the background values for radioactivity that are present under normal circumstances. They can be used as reference values, for instance, during a nuclear emergency.

Radioactivity in air, food, milk, grass and feed

Radioactivity levels in the air were normal, i.e. within the range of previous years. Radioactivity levels in food and milk were well below the export and consumption limits set by European legislation. Radioactivity levels in grass and feed were normal, i.e. within the range of previous years.

Radioactivity in surface water, seawater and drinking water

Radioactivity levels in surface water and seawater were within the range of previous years. Radioactivity levels in untreated water for drinking water production were well below the screening levels above which further investigation should be carried out, with the exception of 19 samples of untreated water (5% of the total number of samples), which were slightly elevated. These measured radioactivity levels do not pose a threat to public health. Further investigation revealed that radioactivity levels in associated finished drinking water were well below the screening levels.

Keywords: radioactivity, environment, airborne particles, water, food, milk

Publiekssamenvatting

Radioactiviteit in het Nederlandse milieu

Resultaten in 2016

In 2016 voldeed Nederland aan de Europese verplichting om jaarlijks de hoeveelheid radioactiviteit in het milieu en in voeding te meten. Alle lidstaten van de Europese Unie zijn volgens het Euratom-verdrag uit 1957 verplicht deze metingen jaarlijks te verrichten. Nederland volgt daarbij de aanbevelingen die in 2000 zijn opgesteld om de metingen volgens een bepaald stramien uit te voeren. De metingen leveren achtergrondwaarden op, ofwel radioactiviteitsniveaus, die onder normale omstandigheden aanwezig zijn. Deze waarden kunnen bij bijvoorbeeld calamiteiten of rampen als referentie dienen. Het RIVM rapporteert namens Nederland aan de Europese Unie over radioactiviteit in het milieu.

Radioactiviteit in lucht, voedsel, melk, gras en veevoer

De radioactiviteitsniveaus in lucht laten een normaal beeld zien dat niet verschilt van voorgaande jaren. De niveaus in voedsel en melk liggen net als in voorgaande jaren duidelijk onder de Europese limieten die zijn opgesteld voor consumptie en export. Ook de radioactiviteitsniveaus in gras en veevoer laten een normaal beeld zien dat niet verschilt van voorgaande jaren.

Radioactiviteit in oppervlaktewater, zeewater en drinkwater

De radioactiviteitsniveaus in oppervlaktewater en zeewater verschillen niet van voorgaande jaren. In ongezuiverd water voor de drinkwaterproductie liggen de niveaus meestal onder de zogeheten screeningswaarden (boven deze waarden moet nader onderzoek worden uitgevoerd). Een uitzondering daarop zijn 19 monsters ongezuiverd water (5 procent van het totale aantal monsters) waarbij licht verhoogde niveaus zijn gemeten. Deze verhogingen zijn zodanig laag dat ze niet schadelijk zijn voor de gezondheid. Aanvullend onderzoek toonde aan dat de niveaus in het gezuiverde drinkwater ruim onder de screeningswaarden lagen.

Trefwoorden: radioactiviteit, milieu, luchtstof, water, voedsel, melk

Acknowledgements

The following institutions contributed to this report:

**The National Institute for Public Health and the Environment
Rijksinstituut voor Volksgezondheid en Milieu (RIVM)**

Data on air dust, deposition, ambient dose rates and drinking water.
Dr. C.P. Tanzi (editor), ing. G.J. Knetsch, ing. R.B. Tax, ing. M. Boshuis,
ir. J.F.M. Versteegh

Rijkswaterstaat (RWS)

Data on seawater and surface water from the main inland water bodies.
C. Engeler, A. Baak

**The Netherlands Food and Consumer Product Safety Authority
Nederlandse Voedsel en Waren Autoriteit (NVWA)**

Data on foodstuffs.
E. Laurensse

RIKILT Wageningen UR

Data on milk and foodstuffs.
ir. S.T. van Tuinen, ing. C. Onstenk, ing. A. Vos van Avezathe

N.V. Elektriciteits-Produktie­maatschappij Zuid-Nederland (EPZ)

Data on environmental samples taken near the Borssele nuclear power
plant, measured by the Nuclear Research & Consultancy Group (NRG).
G.J.L. Goulooze

Contents

Summary — 11

Samenvatting — 13

1 Introduction — 19

2 Airborne particles — 21

- 2.1 Introduction — 21
- 2.2 Results for long-lived α and β activity — 22
- 2.3 Results for γ -emitting radionuclides — 25

3 Deposition — 29

- 3.1 Introduction — 29
- 3.2 Results for long-lived α and β activity — 29
- 3.3 Results for γ -emitting radionuclides — 34

4 National Radioactivity Monitoring Network — 37

- 4.1 Introduction — 37
- 4.2 Results — 38

5 Surface water and seawater — 43

- 5.1 Introduction — 43
- 5.2 The results for surface water — 47
- 5.3 The results for seawater — 58

6 Water for human consumption — 67

- 6.1 Introduction — 67
- 6.2 Results — 67

7 Milk — 69

- 7.1 Introduction — 69
- 7.2 Results — 69

8 Food — 71

- 8.1 Introduction — 71
- 8.2 Results for honey — 72
- 8.3 Results for vegetables and mushrooms — 72
- 8.4 Results for game and poultry — 72
- 8.5 Results for average daily intake — 72

9 Grass & feed — 75

10 The Borssele nuclear power plant — 77

- 10.2 Air — 78
- 10.3 Soil and grass — 79
- 10.4 Water — 80

11 Conclusions — 83

Appendix A Tables of results — 85

Appendix B Presentation of data — 109

Appendix C Glossary — 111

References — 113

Erratum d.d. 24-09-2019 — 108

Summary

The Dutch government is obliged to establish the necessary arrangements to carry out continuous monitoring of the level of radioactivity in air, water and soil under the terms of the Euratom Treaty of 1957. In 2000, the European Commission issued recommendations stipulating the matrices to be measured (air dust, ambient dose, surface water, drinking water, milk and food) and stipulating the frequency of the measurements. The results shall be reported to the European Commission annually.

In 2016, the Netherlands complied with the Euratom recommendations on annually measuring radioactivity in the environment and in food. The measurements were carried out by RIVM, RWS, RIKILT, NVWA and (commissioned by N.V. EPZ) NRG. This report presents the results of radioactivity measurements performed in the Dutch environment in 2016.

Yearly average activity concentrations in air dust were determined for gross α , gross β , ^7Be , ^{137}Cs and ^{210}Pb . The yearly total activity in deposition was determined for gross α , gross β , ^3H , ^7Be , ^{137}Cs , ^{210}Pb and ^{210}Po . Gross α and gross β are the total activity of radionuclides emitting α and β radiation, respectively. The results, which are presented in Table S1 below, are within the range of those presented in previous years.

The National Radioactivity Monitoring Network (NMR) was also used to determine the activity concentrations of gross α and artificial β (β radiation emitted by man-made radionuclides) in air dust. There is a difference between the NMR data and the gross α and gross β data mentioned above, which is due to the contribution of short-lived natural radionuclides (radon daughters) to the NMR data. The yearly average gross α activity concentration in air dust was $3.7 \text{ Bq}\cdot\text{m}^{-3}$. The yearly average of the β activity concentration did not deviate significantly from zero. The NMR was also used to determine the ambient dose equivalent rate: the yearly average was $81 \text{ nSv}\cdot\text{h}^{-1}$.

In surface water, the yearly average activity concentrations of gross α , residual β (gross β minus naturally occurring ^{40}K), ^3H , ^{90}Sr and ^{226}Ra were determined. The yearly average activity concentrations of ^{60}Co , ^{131}I , ^{137}Cs and ^{210}Pb in suspended solids in surface water were also determined. In seawater, the yearly average activity concentrations were determined for gross α , residual β , ^3H and ^{90}Sr . The yearly average activity concentrations of ^{137}Cs and ^{210}Pb in suspended solids in seawater were also determined. The results are presented in Table S1.

The yearly average gross α , residual β , ^3H , ^{90}Sr and ^{226}Ra activity concentrations in surface water were within the range of those found in previous years, with the exception of an elevated concentration of Ra-226 in the Scheldt. The yearly average ^{60}Co , ^{131}I , ^{137}Cs and ^{210}Pb activity concentrations in suspended solids in surface water were within the range of those found in previous years.

The yearly average gross α , residual β , ^3H and ^{90}Sr activity concentrations in seawater were within the range of those found in previous years. The yearly average ^{137}Cs and ^{210}Pb activity concentrations in suspended solids in seawater were within the range of those found in previous years.

Typical activities found in untreated and treated water used for drinking water production are presented in Table S1. There is little potassium (and thus ^{40}K) present in this water. The gross α activity concentration in untreated water used to produce drinking water averaged per production station exceeded the screening level ($0.1 \text{ Bq}\cdot\text{L}^{-1}$) at 6 of the 183 production stations (in 19 of the 388 analyses). Further investigation into these slightly elevated levels in untreated water revealed that the gross α activity concentration in associated finished drinking water was well below the screening level. The gross β activity concentrations were below the screening level ($1.0 \text{ Bq}\cdot\text{L}^{-1}$) and the ^3H activity concentrations were below the parametric value of $100 \text{ Bq}\cdot\text{L}^{-1}$.

The results of the monitoring programme for milk and food are presented in Table S1. Radioactivity was measured in over 600 milk samples and over 2,000 food products. Of these food products, 24 samples of game and poultry and three samples of dried mushrooms contained ^{137}Cs . The set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$ (or $370 \text{ Bq}\cdot\text{kg}^{-1}$ for milk and dairy products) for the activity of radiocesium (sum of ^{134}Cs and ^{137}Cs) was not exceeded.

The measured concentrations of ^{90}Sr , ^{134}Cs and ^{137}Cs in food in $\text{Bq}\cdot\text{kg}^{-1}$ were converted to an average daily intake value per person per day ($\text{Bq}\cdot\text{day}^{-1}$) using food consumption patterns. The average daily intake per person of ^{134}Cs , ^{137}Cs and ^{90}Sr is < 5 , < 7 , $< 5 \text{ Bq}\cdot\text{day}^{-1}$, respectively. The contribution to the effective yearly dose calculated from these average daily intake values is $< 0.1 \text{ mSv}$. The actual daily intake (and following dose contribution) is most likely much lower.

None of the grass and feed samples contained measurable levels of artificial radionuclides (^{60}Co , ^{131}I , ^{132}Te , ^{134}Cs and ^{137}Cs).

Data on environmental samples taken near the Borssele nuclear power plant are presented in Table S2.

Samenvatting

In het kader van het Euratom Verdrag uit 1957 is de Nederlandse overheid verplicht om voorzieningen te treffen om radioactiviteitsgehalten te monitoren in de compartimenten lucht, water en bodem. In 2000 heeft de Europese Commissie aanbevelingen uitgegeven waarin in detail is beschreven wat moet worden gemeten (luchtstof, het omgevingsdosisequivalenttempo, oppervlaktewater, drinkwater, melk en voedsel) en met welke frequentie. De resultaten dienen jaarlijks te worden gerapporteerd aan de Europese Commissie.

Dit rapport bevat de resultaten van radioactiviteitsmetingen in het Nederlandse milieu in 2016. De metingen zijn verricht door RIVM, RWS, RIKILT, NVWA en (in opdracht van N.V. EPZ) NRG. Nederland voldeed in 2016 aan de Europese aanbevelingen ten aanzien van de jaarlijkse radioactiviteitsmetingen in het milieu en in voedsel.

In luchtstof werd de jaargemiddelde activiteitsconcentratie bepaald van totaal- α , totaal- β , ^7Be , ^{137}Cs en ^{210}Pb . In depositie werd de totale jaarlijkse activiteit bepaald van totaal- α , totaal- β , ^3H , ^7Be , ^{137}Cs , ^{210}Pb en ^{210}Po . Totaal- α respectievelijk totaal- β is de totale activiteit aan α - dan wel β -straling uitzendende radionucliden. De resultaten zijn weergegeven in Tabel S1 en vallen binnen de spreiding van de resultaten van voorgaande jaren.

Met het Nationaal Meetnet Radioactiviteit (NMR) werden activiteitsconcentraties bepaald in luchtstof voor totaal- α en kunstmatige β (β -straling uitgezonden door radionucliden, ontstaan door menselijk handelen). Er is een verschil tussen de NMR-metingen en bovenstaande totaal- α - en totaal- β -metingen, wat wordt veroorzaakt door de bijdrage van kortlevende natuurlijke radionucliden (radondochters). Het jaargemiddelde voor de totaal- α -activiteitsconcentratie in luchtstof was $3,7 \text{ Bq}\cdot\text{m}^{-3}$. Het jaargemiddelde voor de kunstmatige β -activiteitsconcentratie in luchtstof week niet significant af van nul. Met het NMR werd daarnaast het omgevingsdosisequivalenttempo bepaald; het jaargemiddelde was $81 \text{ nSv}\cdot\text{h}^{-1}$.

In oppervlaktewater werd de jaargemiddelde activiteitsconcentratie bepaald van totaal- α , rest- β (totaal- β minus het van nature aanwezige ^{40}K), ^3H , ^{90}Sr en ^{226}Ra en de jaargemiddelde activiteitsconcentratie van ^{60}Co , ^{131}I , ^{137}Cs en ^{210}Pb in zwevend stof.

In zeewater werd de jaargemiddelde activiteitsconcentratie bepaald van totaal- α , rest- β , ^3H en ^{90}Sr . In zwevend stof in zeewater werd de jaargemiddelde activiteitsconcentratie bepaald van ^{137}Cs en ^{210}Pb . De resultaten zijn weergegeven in Tabel S1.

De jaargemiddelde activiteitsconcentraties van totaal- α , rest- β (totaal- β minus het van nature aanwezige ^{40}K), ^3H , ^{90}Sr en ^{226}Ra in oppervlaktewater vallen binnen de spreiding van de resultaten van voorgaande jaren, met de uitzondering van verhoogde concentratie ^{226}Ra in de Schelde. De jaargemiddelde activiteitsconcentraties van ^{60}Co ,

^{131}I , ^{137}Cs en ^{210}Pb in zwevend stof in oppervlaktewater vallen binnen de spreiding van de resultaten van voorgaande jaren.

De jaargemiddelde totaal α -, rest β -, ^3H - en ^{90}Sr -activiteitsconcentraties in zeewater vallen binnen de spreiding van de resultaten van voorgaande jaren. De jaargemiddelde ^{137}Cs - en ^{210}Pb -activiteitsconcentraties in zwevend stof in zeewater vallen binnen de spreiding van de resultaten van voorgaande jaren.

Gangbare activiteitsconcentraties die in ongezuiverd en gezuiverd water voor de drinkwaterproductie gevonden worden, zijn weergegeven in Tabel S1. In dit water is een geringe hoeveelheid kalium, en dus ^{40}K , aanwezig. In 2016 overschrijdt de totaal α -activiteitsconcentratie in ongezuiverd water voor drinkwaterproductie per productiestation de screeningswaarde van $0,1 \text{ Bq}\cdot\text{L}^{-1}$ bij 6 van de 183 productiestations (in 19 van de 388 uitgevoerde analyses). Deze verhogingen zijn zodanig laag dat ze niet schadelijk zijn voor de gezondheid. Additioneel onderzoek naar aanleiding van de licht verhoogde niveaus in ongezuiverd water toonde aan dat de niveaus in het gezuiverde drinkwater ruim onder de screeningswaarden lagen. De totaal β -activiteitsconcentraties waren lager dan $1,0 \text{ Bq}\cdot\text{L}^{-1}$ en de ^3H -activiteitsconcentraties waren lager dan $100 \text{ Bq}\cdot\text{L}^{-1}$.

De resultaten van het meetprogramma voor melk en voedsel zijn weergegeven in Tabel S1. Radioactiviteit werd geanalyseerd in ruim 600 melkmonsters en meer dan 2000 voedselproducten, waarvan 24 monsters pluimvee en wild en drie monsters gedroogde paddenstoelen ^{137}Cs bevatten. De limiet van $600 \text{ Bq}\cdot\text{kg}^{-1}$ (respectievelijk $370 \text{ Bq}\cdot\text{kg}^{-1}$ voor melk en melkproducten) van radiocesium (som van ^{134}Cs en ^{137}Cs) wordt niet overschreden.

De gemeten concentraties ^{90}Sr , ^{134}Cs en ^{137}Cs ($\text{Bq}\cdot\text{kg}^{-1}$) in voedsel worden omgerekend naar een gemiddelde dagelijkse opname per persoon per dag ($\text{Bq}\cdot\text{dag}^{-1}$) door gebruik te maken van voedselconsumptiepatronen. De gemiddelde dagelijkse opname per persoon is < 5 , < 7 , $< 5 \text{ Bq}\cdot\text{dag}^{-1}$ voor respectievelijk ^{134}Cs , ^{137}Cs en ^{90}Sr . De bijdrage aan de effectieve jaardosis berekend uit deze waarden is $< 0,1 \text{ mSv}$. De werkelijke dagelijkse opname (en daaruit volgende bijdrage aan de jaardosis) is vermoedelijk veel lager.

Geen van de geanalyseerde monsters gras en veevoer bevatten meetbare hoeveelheden van kunstmatige radionucliden (^{60}Co , ^{131}I , ^{132}Te , ^{134}Cs en ^{137}Cs).

Gegevens betreffende milieumonsters genomen rondom de kerncentrale Borssele zijn weergegeven in Tabel S2.

Table S1 Summary of the results from the Dutch monitoring programme in 2016

Matrix	Parameter	Locations	Values	Frequency (per year)
Air dust ⁽¹⁾	Gross α	1	0.019 mBq·m ⁻³	52
	Gross β	1	0.362 mBq·m ⁻³	52
	⁷ Be	1	2.910 mBq·m ⁻³	52
	¹³⁷ Cs	1	0.000208 mBq·m ⁻³	52
	²¹⁰ Pb	1	0.312 mBq·m ⁻³	52
Deposition ⁽²⁾	Gross α	1	66 Bq·m ⁻²	12
	Gross β	1	88 Bq·m ⁻²	12
	³ H ⁽³⁾	1	22–1,510 Bq·m ⁻²	12
	⁷ Be	1	1,375 Bq·m ⁻²	52
	¹³⁷ Cs ⁽³⁾	1	0.04–1.15 Bq·m ⁻²	52
	²¹⁰ Pb	1	95–98 Bq·m ⁻²	52
	²¹⁰ Po	1	40.6 Bq·m ⁻²	12
Surface water ⁽¹⁾	Gross α	8	29–249 mBq·L ⁻¹	12–13 ⁽⁴⁾
	Residual β	8	13–112 mBq·L ⁻¹	12–13 ⁽⁴⁾
	³ H	8	1,550–14,600 mBq·L ⁻¹	6–13 ⁽⁴⁾
	⁹⁰ Sr	3	< 1–3.1 mBq·L ⁻¹	6–7 ⁽⁴⁾
	²²⁶ Ra	4	4.0–33.1 mBq·L ⁻¹	6–7 ⁽⁴⁾
Suspended solids in surface water ⁽¹⁾	⁶⁰ Co	8	< 1–10.4 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	¹³¹ I	8	< 1–36.0 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	¹³⁷ Cs	8	3.0–12.4 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	²¹⁰ Pb	4	80–140 Bq·kg ⁻¹	6–7 ⁽⁴⁾
Seawater ⁽¹⁾	Gross α	8	190–420 mBq·L ⁻¹	4–13 ⁽⁴⁾
	Residual β	8	34–140 mBq·L ⁻¹	4–13 ⁽⁴⁾
	³ H	8	350–5,200 mBq·L ⁻¹	4–13 ⁽⁴⁾
	⁹⁰ Sr	4	< 1.4–1.9 mBq·L ⁻¹	4–13 ⁽⁴⁾
Suspended solids in seawater ⁽¹⁾	¹³⁷ Cs	1	3.3 Bq·kg ⁻¹	4 ⁽⁴⁾
	²¹⁰ Pb	1	62 Bq·kg ⁻¹	4 ⁽⁴⁾
Drinking water ⁽¹⁾	Gross α	183	< 0.06 Bq·L ⁻¹	388 ⁽⁵⁾
	Gross β	186	< 0.1 Bq·L ⁻¹	428 ⁽⁵⁾
	Residual β	168	< 0.1 Bq·L ⁻¹	390 ⁽⁵⁾
	³ H	88	< 4.0 Bq·L ⁻¹	448 ⁽⁵⁾
Milk ⁽¹⁾	⁴⁰ K	22	51.2 Bq·kg ⁻¹ ⁽⁶⁾ 50.1 Bq·kg ⁻¹ ⁽⁷⁾	581 ⁽⁵⁾ 11
	⁶⁰ Co	22	< 1.4 Bq·kg ⁻¹	581 ⁽⁵⁾
	⁹⁰ Sr	22	< 5 Bq·kg ⁻¹	47 ⁽⁵⁾
	¹³¹ I	22	< 0.6 Bq·kg ⁻¹	581 ⁽⁵⁾
	¹³⁴ Cs	22	< 0.6 Bq·kg ⁻¹	581 ⁽⁵⁾
	¹³⁷ Cs	22	< 0.5 Bq·kg ⁻¹	581 ⁽⁵⁾

Continued on next page

Table S1 Continued

Matrix	Parameter	Locations	Values	Frequency (per year)
Food (8, 9)				
Grain and grain products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	31 (0) (11)
Vegetables and mushrooms	¹³⁷ Cs (10)	-	22–1019 Bq·kg ⁻¹	84 (3) (11, 12)
Fruit and fruit products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	10 (0) (11)
Milk and dairy products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	50 (0) (11)
Meat and meat products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	32 (0) (11)
Game and poultry	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	26 (0) (11)
Salads	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	17 (0) (11)
Oil and butter	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	30 (0) (11)
Honey	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	49 (1) (11)
Tea	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	3 (0) (11)
Mineral water	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	18 (0) (11)
Fish	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	21 (0) (11)
Food (8, 13)				
Vegetables and fruits	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	287 (0) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	23 (0) (11)
Meat and meat products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	568 (0) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	15 (0) (11)
Game and poultry	¹³⁷ Cs (10)	-	5–43 Bq·kg ⁻¹	398 (24) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	19 (0) (11)
Eggs	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	156 (0) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	6 (0) (11)
Fish and seafood products	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	201 (0) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	24 (0) (11)
Ready meals	¹³⁷ Cs (10)	-	< 5 Bq·kg ⁻¹	45 (0) (11)
	⁹⁰ Sr	-	< 5 Bq·kg ⁻¹	45 (0) (11)

(1) Yearly average.

(2) Yearly total.

(3) A 68% confidence interval.

(4) Frequency depends on location.

(5) Total number of samples taken combined over all locations.

(6) Yearly average in cow's milk.

(7) Yearly average in goat's milk.

(8) Given range represents values of individual (positive) samples.

(9) As measured by the Netherlands Food and Consumer Product Safety Authority.

(10) Samples were analysed for ¹³⁴Cs as well, but measurements were below the detection limit of 5 Bq·kg⁻¹.

(11) Total number of samples taken (number of samples where the minimum detectable activity is exceeded in brackets).

(12) Activity concentrations above 1,000 Bq·kg⁻¹ were measured in dried mushrooms. As mushrooms are not consumed in their dry form, the activity concentration in their consumable form is compared to the set limit of 600 Bq·kg⁻¹.

(13) As measured by RIKILT Wageningen UR.

Table S2 Summary of the results of the monitoring programme in the vicinity of the Borssele nuclear power plant in 2016

Matrix	Parameter	Locations	Values ⁽¹⁾	Frequency (per year)
Air (dust)	Gross α	5	0.002–0.153 mBq·m ⁻³	12
	Gross β	5	0.03–1.27 mBq·m ⁻³	12
	⁶⁰ Co	5 ⁽²⁾	< 0.04–< 0.08 mBq·m ⁻³	12
	¹³¹ I _{el} ⁽³⁾	5 ⁽²⁾	< 0.1–< 0.3 mBq·m ⁻³	12
	¹³¹ I _{or} ⁽⁴⁾	5 ⁽²⁾	< 0.4–< 1 mBq·m ⁻³	12
	¹³⁷ Cs	5 ⁽²⁾	< 0.03–< 0.06 mBq·m ⁻³	12
	Nat. ⁽⁵⁾	5 ⁽²⁾	< 1.3–3 mBq·m ⁻³	12
Grass	⁶⁰ Co	5 ⁽²⁾	< 0.8–< 3 Bq·kg ⁻¹	12
	¹³¹ I	5 ⁽²⁾	< 0.9–< 2 Bq·kg ⁻¹	12
	¹³⁷ Cs	5 ⁽²⁾	< 0.7–< 2 Bq·kg ⁻¹	12
Soil	⁵⁴ Mn	4	< 0.3–< 0.4 Bq·kg ⁻¹	1
	⁶⁰ Co	4	< 0.3–< 0.4 Bq·kg ⁻¹	1
	¹³⁴ Cs	4	< 0.2–< 0.4 Bq·kg ⁻¹	1
	¹³⁷ Cs	4	1.27–1.48 Bq·kg ⁻¹	1
Water	Residual β	4	< 0.011–0.132 Bq·L ⁻¹	12
	³ H	4	1.4–7.3 Bq·L ⁻¹	12
Suspended solids	Gross β	4	0.20–3.10 kBq·kg ⁻¹	12
Seaweed	⁶⁰ Co	4 ⁽²⁾	< 0.6–< 2 Bq·kg ⁻¹	12
	¹³¹ I	4 ⁽²⁾	< 0.5–< 2 Bq·kg ⁻¹	12
	¹³⁷ Cs	4 ⁽²⁾	< 0.5–< 2 Bq·kg ⁻¹	12
Sediment	⁶⁰ Co	4 ⁽²⁾	< 0.2–< 0.3 Bq·kg ⁻¹	12
	¹³¹ I	4 ⁽²⁾	< 0.2–< 0.3 Bq·kg ⁻¹	12
	¹³⁷ Cs	4 ⁽²⁾	< 0.2–0.99 Bq·kg ⁻¹	12

⁽¹⁾ Given range represents the range of values of individual samples.

⁽²⁾ Analysis was performed on a combined sample of the monthly samples collected in different locations.

⁽³⁾ Elemental ¹³¹I.

⁽⁴⁾ Organically bound ¹³¹I.

⁽⁵⁾ Naturally occurring γ -emitters.

1 Introduction

Radioactivity of natural and artificial origin can be found in low concentration levels all around the globe. So-called naturally occurring radionuclides can be found in soil, water and air, and have existed since the creation of the universe or are continually generated, either from the decay of primordial isotopes or as a result of cosmic radiation. Concentration levels of naturally occurring radionuclides, such as ^{40}K and daughters from the uranium and thorium series, may be enhanced as a result of human activities within or outside national borders (e.g. discharges from the ore-processing industry).

Man-made radionuclides are found in the environment as a result of events such as nuclear weapons tests and discharges from medical facilities and nuclear installations.

It is important to monitor radioactivity levels in the environment, as the intake of radioactivity and exposure to ionising radiation can lead to an effective dose that may eventually result in adverse health effects. Monitoring radioactivity levels in the environment provides background radioactivity levels under normal circumstances that enable the detection and confirmation of abnormal levels.

This report presents the results of radioactivity measurements performed in the environment in the Netherlands in 2016.

The aim of this report is threefold:

- i) to present a survey of radioactivity measurements performed in the Dutch environment, providing information on the exposure of the population to ionising radiation;
- ii) to provide information on typical environmental radioactivity levels, which can be used as a reference in the event of a radiological or nuclear incident or emergency;
- iii) to show the compliance of monitoring programmes in the Netherlands with the corresponding European obligations and recommendations and to report possible omissions.

In the following chapters, the results of the measurements will be presented in graphs and tables. More detailed information is presented in Appendix A. Chapters 2 to 8 are subdivided according to the structure of the Commission Recommendation on the Application of Article 36 of the Euratom Treaty [1] and present the results of measurements for various environmental compartments. Chapter 9 contains data on radioactivity levels in grass and feed. Chapter 10 contains data on environmental samples taken near the Borssele nuclear power plant. General conclusions are presented in Chapter 11.

Appendix B describes the methods used for the presentation of data. A glossary of frequently occurring terms is given in Appendix C.

2 Airborne particles

2.1 Introduction

Table 2.1 describes the monitoring programme for the determination of radionuclides in air dust. The sampling was performed at the RIVM premises in Bilthoven, the Netherlands. Air dust samples for the measurement of gross α , gross β and γ -emitters were collected weekly using a high-volume sampler. The high-volume sampler described in [2] was replaced by a Snow White high-volume sampler from Senya Ltd [3] in 2011.

The change in equipment coincided with a change in the filter type (polypropylene G-3 instead of glass fibre GF10), the volume sampled (125,000 m³ instead of 50,000 m³) and the sampling height (on top of a three-storey building instead of 1.8 m above ground level). Samples were collected weekly according to a standard procedure [4].

The collection efficiency of the filter type G-3 was determined to be $96 \pm 1\%$ with a flow rate of approximately 760 (normalized) m³·h⁻¹ based on ⁷Be and ²¹⁰Pb results [3]. The results presented in this chapter take this collection efficiency into account.

After sampling, the G-3 filters were dried and weighed to determine the dust load. Then a sub-sample was taken from the filter for the determination of gross α and gross β according to a standard procedure [5]. The remainder of the filter was folded into a 250 ml container and measured on a coaxial detector (4 days delay time, 100,000 seconds counting time) to determine volatile γ -emitters according to standard procedures [5, 6].

Following this measurement, the filter was dry-ashed at 450 °C for 16 h. Calcium sulphate was added to the resulting residue to achieve a sample of 4 g, which was homogenized and transferred to a polyethylene vial. Measurements were carried out on a coaxial well-type detector (178,200 seconds counting time) according to standard procedures [5, 6].

The period between sampling and the gross α and gross β analysis was five to ten days, which is long compared with the decay time of the short-lived decay products ²²²Rn and ²²⁰Rn. This is done to ensure that these naturally occurring short-lived decay products do not contribute to the measured α and β activity concentrations.

In 2005 the data from 1991 to 2004 were re-analysed to determine the yearly averages following the method described in Appendix B [7]. This might have led to small differences between the data presented in this report and the data reported prior to 2005.

Table 2.1 Monitoring programme for the determination of radionuclides in air dust

Matrix	Location	Parameter	Sample period	Sample volume	Analysis frequency
Air dust	Bilthoven	gross α , gross β	1 week	925 m ³ ⁽²⁾	weekly
Air dust	Bilthoven	γ -emitters ⁽¹⁾	1 week	125,000 m ³	weekly

⁽¹⁾ γ -spectroscopic analysis of specific γ -emitting radionuclides.

⁽²⁾ A sub-sample of 0.74% from the total filter, through which approximately 125,000 m³ of air was sampled.

2.2 Results for long-lived α and β activity

The weekly results of gross α and β activity concentrations in air dust are given in Figure 2.1 and Table A1 (see Appendix A). Due to large uncertainties caused by variations in the amount of dust on the filters, gross α activity concentrations in air dust should be regarded as indicative values [5]. The frequency distributions of gross α activity and gross β activity concentrations in air dust are given in Figures 2.2 and 2.3, respectively.

The yearly average activity concentrations in air dust were 0.019 mBq·m⁻³ for gross α and 0.362 ± 0.006 mBq·m⁻³ for gross β . The yearly averages of the gross α and β activity concentrations of long-lived radionuclides in 2016 were within the range of the results from the period 1992–2015, as illustrated in Figure 2.4. Since 2007, a new (more realistic) calibration for gross α has been applied to the measurements. The current calibration factor is 1.4 times higher than the one used before 2007, resulting in lower reported gross α activities.

In 2011, a change in equipment, with a coinciding change in filter type, resulted in changes in the reported gross α (-24%) and gross β (-15%) results, for which no correction is applied [3]. A possible explanation for these changes is a deeper permeation of the air dust in the present filter type G-3 than in the previous filter type GF10. This results in a difference in the self-absorption of the α and β particles measured, i.e. lower gross α and gross β results in the present G-3 filter than in the previous GF10 filter.

Normally there is a good correlation between gross β and naturally occurring ²¹⁰Pb (in equilibrium with its β -emitting daughter ²¹⁰Bi) activity concentrations, which suggests that under normal conditions ²¹⁰Pb/²¹⁰Bi is the main contributor to the gross β value (see Figure 2.8).




Figure 2.1 Weekly average gross α and β activity concentrations of long-lived radionuclides in air dust sampled at RIVM




Figure 2.2 Frequency distribution of gross α activity concentration of long-lived radionuclides in air dust collected weekly in 2016




Figure 2.3 Frequency distribution of gross β activity concentration of long-lived radionuclides in air dust collected weekly in 2016




Figure 2.4 Yearly average gross α and gross β activity concentrations of long-lived radionuclides in air dust at RIVM since 1992

2.3 Results for γ -emitting radionuclides

Several γ -emitting radionuclides were detected frequently in air dust: ^7Be (52 times), ^{210}Pb (52 times) and ^{137}Cs (47 times). The results are presented in Table A3 and Figures 2.5, 2.6 and 2.7. The detection limits for radionuclides considered in the γ -spectroscopic analysis of the HVS samples are given in Table A2.

The yearly average activity concentrations of ^7Be , ^{137}Cs and ^{210}Pb were $2,910 \pm 30$, 0.208 ± 0.004 and $312 \pm 4 \mu\text{Bq}\cdot\text{m}^{-3}$, respectively.

The behaviour of ^7Be in the atmosphere has been studied worldwide [8, 9, 10, 11, 12, 13, 14]. Natural ^7Be (half-life of 53.3 days) is formed by spallation reactions of cosmogenic radiation with atmospheric nuclei such as nitrogen and oxygen, resulting in the formation of BeO or $\text{Be}(\text{OH})_2$ molecules. Approximately 70% of ^7Be is produced in the stratosphere and the remainder is produced in the troposphere. It has an estimated residence time of one to two years in the stratosphere and approximately six weeks in the troposphere. Most of the ^7Be produced in the stratosphere does not reach the troposphere, except during spring, when seasonal thinning of the tropopause takes place at mid-latitudes, resulting in air exchange between the stratosphere and the troposphere. In the troposphere, ^7Be rapidly associates mainly with submicron-sized aerosol particles. Gravitational settling and precipitation processes accomplish transfer to the earth's surface. Seasonal variations in the concentration of ^7Be in surface air are influenced by the following main atmospheric processes: wet and dry deposition, mass exchange between stratosphere and troposphere, vertical transport in the troposphere, and horizontal transport of air masses from the subtropics and mid-latitudes to the tropics and polar regions.

The red line in Figure 2.5 shows the seasonal variation of the ^7Be activity concentration, with peaks during the spring and summer periods, reflecting the seasonal variations in the transport rate of air from the stratosphere to the troposphere. Figure 2.5 further shows the influence of the solar cycle. The maxima of 1997 and 2007–2009, and the minimum at 2000–2002 are consistent with the solar minima (measured by radio flux and sunspot count) of 1996–1997 and 2008–2009, and the solar maximum of 2000–2002, respectively [15]. In the summer of 1991, two severe geomagnetic storms caused a significant worldwide disturbance of the earth's geomagnetic field. This resulted in a considerable decrease in cosmogenic radiation, which was unprecedented in at least the previous four decades [16]. The absence of a 1991 summer peak in the ^7Be activity concentration can be explained by the decrease in cosmogenic radiation. The concentrations found for ^7Be in 2016 fit into the pattern described above.




Figure 2.5 Weekly average ^7Be activity concentrations (blue) in air dust at RIVM since 1991. The red line is a moving average of 13 weeks.

The nuclide ^{137}Cs (half-life of 30.2 years) is of anthropogenic origin. The two main sources of ^{137}Cs in the environment are atmospheric nuclear weapons tests and some nuclear accidents, most notably the Chernobyl accident of 1986 and the Fukushima Daiichi accident of 2011. Resuspension of previously deposited activity has been the main source of airborne ^{137}Cs activity in the Netherlands since 1986.

Figure 2.6 shows a peak during May 1992. During that month, several wildfires occurred near the Chernobyl area [17], and the level of airborne ^{137}Cs activity increased ten times in the 30 km exclusion zone around Chernobyl. It is possible that the airborne ^{137}Cs was transported to Western Europe by a strong easterly wind in the same period [18]. On 29 May 1998, an incident occurred at Algeciras (Spain): an iron foundry melted a ^{137}Cs source concealed in scrap metal [19]. As a result, elevated levels of airborne ^{137}Cs activity were measured in France, Germany, Italy and Switzerland during late May and early June. Figure 2.6 shows a slightly elevated level of ^{137}Cs activity (second peak) around the same period (29 May to 5 June 1998). Such slightly elevated levels are not uncommon, as can be seen in Figure 2.6. These elevations may be related to the resuspension of previously deposited dust, especially during a period of strong winds from the continent [19]. From 18 March to 10 June 2011, elevated levels of ^{137}Cs activity were measured as a result of the incident at Fukushima (Japan). More detailed results on ^{137}Cs and other radionuclides during that period are presented in [20].

Figure 2.6 shows a lack of data between 2000 and the middle of 2009. During that period the detection limit was higher than it had been during 1991–1999 due to a different detector set-up. Since July 2009, a new detector set-up has been used, which results in lower detection limits (similar to those used before 2000).




Figure 2.6 Weekly average ^{137}Cs activity concentrations in air dust at RIVM since 1991

The primary source of atmospheric ^{210}Pb (half-life of 22.3 years) is the decay of ^{222}Rn exhaled from continental surfaces. Therefore, the atmospheric concentration of ^{210}Pb over continental areas is generally higher than it is over oceanic areas (^{222}Rn exhalation from the ocean is 1,000 times less than it is from the continents). The reported UNSCEAR reference level of ^{210}Pb in air dust is $500 \mu\text{Bq}\cdot\text{m}^{-3}$ [21]. In the atmosphere, this radionuclide is predominantly associated with submicron-sized aerosol particles [22, 23]. The mean aerosol (carrying ^{210}Pb) residence time in the troposphere is approximately 5 days [24].

Other sources of ^{210}Pb in air dust are volcanic activity and industrial emissions [25, 26, 27, 28, 29, 30]. Examples of industrial emissions are discharges from power plants that use fossil fuels, discharges from the fertilizer and phosphorus industries, and exhaust gases from traffic. In the Netherlands, emissions by power plants are only of local importance with respect to ^{210}Pb deposition. Emissions by the phosphorus industry contribute a negligible part of the yearly total ^{210}Pb deposition [30]. Furthermore, the phosphorus industry ceased to be operational in the Netherlands in 2012. Volcanic eruptions bring uranium decay products into the atmosphere, such as ^{226}Ra , ^{222}Rn , ^{210}Pb and ^{210}Po . Beks et al. [27] estimate that volcanoes contribute $60 \text{ TBq}\cdot\text{year}^{-1}$ to the atmospheric ^{210}Pb stock. Unusual (high) ^{210}Pb values might be explained by natural phenomena such as an explosive volcanic eruption, Saharan dust [31, 32, 33] or the resuspension of (local) dust. Normally there is a good correlation between ^{210}Pb (in equilibrium with its β -emitting daughter ^{210}Bi) and gross β activity concentrations, as was the case in 2016 (Figure 2.8). This suggests that, under normal conditions, $^{210}\text{Pb}/^{210}\text{Bi}$ is the main contributor to the gross β value. The weekly average ^{210}Pb activity concentrations in 2016 were within the range of those found in previous years (Figure 2.7).




Figure 2.7 Weekly average ^{210}Pb activity concentrations in air dust at RIVM since 1991




Figure 2.8 The weekly average gross β and ^{210}Pb activity concentrations in air dust at RIVM show a good correlation.

3 Deposition

3.1 Introduction

Table 3.1 describes the monitoring programme for determining radionuclides in deposition. Sampling was done on the RIVM premises in Bilthoven. Samples were collected weekly for γ -emitters and monthly for gross α , gross β , ^3H and ^{210}Po according to a standard procedure [34].

The samples – weekly for γ -emitters and monthly for gross α and gross β – were acidified with sulphuric acid and evaporated. The resulting sulphate residue was analysed according to standard procedures [6, 35, 36].

The monthly samples for ^3H were made alkaline by the addition of sodium carbonate and then distilled. A 10 ml aliquot of the distillate was mixed with an equal amount of scintillation solution (Ultima Gold LLT) in a plastic counting vial and then counted on a liquid scintillation counter for 1,000 minutes per sample.

The monthly samples for ^{210}Po were reduced in volume by evaporation. The resulting solution was analysed according to a standard procedure [37] with a minor difference: the ingrowth of ^{210}Po from ^{210}Pb was derived from the ^{210}Pb results from the weekly samples for γ -emitters, rather than by using the procedure described in [37].

In 2005 the data from 1993 to 2004 were re-analysed to determine the yearly totals by the method described in Appendix B [7]. This can lead to small differences between the data presented in this report and the data reported prior to 2005.

Table 3.1 Monitoring programme for the determination of radionuclides in deposition


Matrix	Location	Parameter	Sample period	Sample volume	Analysis frequency
Deposition	Bilthoven	γ -emitters ⁽¹⁾	1 week	variable	weekly
Deposition	Bilthoven	gross α , gross β , ^{210}Po	1 month	variable	monthly
Deposition	Bilthoven	^3H	1 month	variable	quarterly

⁽¹⁾ γ spectroscopic analysis of specific γ emitting radionuclides.


3.2 Results for long-lived α and β activity

The monthly deposited gross α and gross β activities of long-lived radionuclides are given in Figure 3.1, Figure 3.3 and Table A4. The yearly total depositions of gross α and gross β were 66 ± 2 and $88 \pm 2 \text{ Bq}\cdot\text{m}^{-2}$, respectively. The yearly total depositions of gross α and gross β are within the range of those from previous years, as illustrated in Figure 3.2, Figure 3.4 and Table A6.

The monthly deposition of ^3H is given in Table A4. In 2016 the yearly total deposition of ^3H ranged between 22 and 1,510 $\text{Bq}\cdot\text{m}^{-2}$ (68% confidence interval). The yearly total consisted of 12 samples and 11 out of 12 measurements were below the detection limit. These detection limits were used for the contribution to the yearly total, as described in Appendix B. The range in 2016 did not differ significantly from those measured since 1993, as illustrated in Figure 3.5 and Table A6. Up to 1998, samples were electrolytically enriched before counting, which resulted in a much lower detection limit than the one used from 1998 onwards.




Monthly totals (black dots) are shown with a 68% confidence interval (coloured bars).
Figure 3.1 Monthly deposition of gross alpha activity of long-lived radionuclides at RIVM




Yearly totals (black dots) are shown with a 68% confidence interval (coloured bars). Only the 68% confidence interval is shown if the sum leading to the yearly total contains at least one detection limit.

Figure 3.2 Yearly deposition of gross α activity of long-lived radionuclides at RIVM since 1993




Monthly totals (black dots) are shown with a 68% confidence interval (coloured bars).


Figure 3.3 Monthly deposition of gross β activity of long-lived radionuclides at RIVM



Yearly totals (black dots) are shown with a 68% confidence interval (coloured bars).
 Figure 3.4 Yearly deposition of gross β activity of long-lived radionuclides at RIVM since 1993




Yearly totals (black dots) are shown with a 68% confidence interval (coloured bars). Only the 68% confidence interval is shown if the sum leading to the yearly total contains at least one detection limit.
 Figure 3.5 Yearly deposition of ^3H at RIVM since 1993



Monthly totals (black dots) are shown with a 68% confidence interval (coloured bars). The result for November was rejected.

Figure 3.6 Monthly deposition of ^{210}Po activity at RIVM



Yearly totals (black dots) are shown with a 68% confidence interval (coloured bars). Only the 68% confidence interval is shown if the sum leading to the yearly total contains at least one detection limit.


Figure 3.7 Yearly deposition of ^{210}Po activity at RIVM since 1993

The monthly a spectroscopy results for ^{210}Po are given in Figure 3.6 and Table A5. The results for previous years are given in Figure 3.7 and Table A7. The yearly total deposition of ^{210}Po in 2016 was $40.6 \pm 1.0 \text{ Bq}\cdot\text{m}^{-2}$ (68% confidence interval) and is based on 11 monthly results. This value is within the range of the values from previous years, as illustrated in Figure 3.7 and Table A7.


3.3 Results for γ -emitting radionuclides

The naturally occurring radionuclides ^7Be and ^{210}Pb were found in all 52 and in 51 of the 52 weekly deposition samples, respectively. The yearly total deposition of ^7Be was $1,375 \pm 18 \text{ Bq}\cdot\text{m}^{-2}$ and the yearly total deposition of ^{210}Pb ranged between 95 and $98 \text{ Bq}\cdot\text{m}^{-2}$. The nuclide ^{137}Cs was detected in two of the 52 weekly samples (the detection limit for ^{137}Cs is $0.02 \text{ Bq}\cdot\text{m}^{-2}$). The yearly total deposition of ^{137}Cs ranged between 0.04 and $1.15 \text{ Bq}\cdot\text{m}^{-2}$ (68% confidence interval). The weekly results for deposition of ^7Be , ^{137}Cs and ^{210}Pb are given in Table A8 and Figures 3.9 and 3.12. The results for previous years are given in Table A7 and Figures 3.10, 3.11 and 3.13.

Figure 3.11 shows a change in trend for ^{137}Cs between 2000 and the middle of 2009. During that period the detection limit was higher than it had been during 1993–1999 due to a different detector set-up. Since July 2009, a new detector set-up has been used, which results in lower detection limits. Since the ^{137}Cs deposition values are mostly below the detection limit, no conclusion can be drawn concerning the correlation between the measured ^{137}Cs values in air dust and the measured ^{137}Cs deposition values.




Weekly totals (black dots) are shown with a 68% confidence interval (coloured bars).
 Figure 3.9 Weekly deposited ^7Be activity at RIVM




Yearly totals (black dots) are shown with a 68% confidence interval (coloured bars). Only the 68% confidence interval is shown if the sum leading to the yearly total contains at least one detection limit.

Figure 3.10 Yearly ⁷Be activity deposited at RIVM since 1993




Yearly averages are shown solely as a 68% confidence interval since the sum leading to the yearly total contains at least one detection limit.

Figure 3.11 Yearly ¹³⁷Cs activity deposited at RIVM since 1993



Weekly averages (black dots) are shown with a 68% confidence interval (coloured bars). In week 49 the measurement was below the detection limit of 0.2 Bq·m⁻² (not shown).

Figure 3.12 Weekly deposited ²¹⁰Pb activity at RIVM



Yearly averages (black dots) are shown with a 68% confidence interval (coloured bars). Only the 68% confidence interval is shown if the sum leading to the yearly total contains at least one detection limit.

Figure 3.13 Yearly ²¹⁰Pb activity deposited at RIVM since 1993

4 National Radioactivity Monitoring Network

4.1 Introduction

This chapter presents data on gross α and artificial β activity concentrations in air dust and ambient dose equivalent rates, as measured by the National Radioactivity Monitoring Network (Nationaal Meetnet Radioactiviteit, NMR). The data on gross α and artificial β differ in sample size, sampling frequency and analytical procedures (including a different approach to the contribution of short-lived natural radionuclides, i.e. radon daughters) from those given in Chapter 2.

The NMR consists of 167 sites, at which the ambient dose equivalent rate is determined. At 14 measuring sites, gross α and artificial β activity concentrations are determined, as well as the ambient dose equivalent rate (at a height of 3.5 m above ground level) [38]. At the other 153 measuring sites, only the ambient dose equivalent rate is determined (at 1 m above ground level).

Since the dose equivalent rate monitors are placed differently at 14 of the 167 sites with regard to height and surface covering, results can differ between the two types of measuring site [39]. For this reason, the 14 dose equivalent rate monitors are not taken into account when calculating the yearly average ambient dose equivalent. The reported artificial β activity concentrations are calculated from the difference between the measured gross β activity concentration and the natural gross β activity derived from the measured gross α activity concentration.

During the second half of 2002, the 14 aerosol FAG FHT59S monitors were gradually replaced by 14 new Berthold BAI 9128 monitors. Due to differences in detection method, filter transport, calibration radionuclides and algorithms, the results for the activity concentrations from the two types of monitor are not exactly the same. By running both monitors simultaneously at the same location, the measured gross α activity concentration was compared. On average, the Berthold monitor systematically reported about 20% higher values than the FAG monitor [40]. The estimated random uncertainty for both types of monitor is about 20%. No correction was applied for the difference in the gross α activity concentration between the Berthold and FAG monitors.

From November 2014 until the end of 2015 most of the ambient dose equivalent monitors were replaced. The Bitt RS03 monitors (proportional counters) were replaced by new Saphymo XL-2-3 monitors (Geiger-Müller). The energy response, cosmic response and self-effect of the two types of monitor differ slightly. Compared with the Bitt monitor, the Saphymo monitor reports on average 8 nSv/h higher at the natural background radiation level in the Netherlands. No correction for this difference is applied.

The data presented in this chapter are based on 10-minute measurements. Averages over the year are calculated for each location,

using daily averages derived from the 10-minute measurements (Tables A9 and A10). The data on external radiation, expressed in ambient dose equivalents, contain a systematic uncertainty because of an overestimation of the cosmogenic dose rate. NMR data are not corrected for these response uncertainties.

4.2 Results


An impression of the spatial variation in the yearly averages of the NMR data, constructed by using the RIVM's Geographical Information System (GIS), is shown in Figures 4.1 and 4.3. An inverse distance weight interpolation algorithm was applied to calculate values between the NMR stations.

Figure 4.2 presents the yearly averages of gross α activity concentration since 1990, while Figure 4.4 presents the yearly averages of ambient dose equivalent rate since 1996.

In 2016 the yearly average gross α activity concentration in air dust was $3.7 \text{ Bq}\cdot\text{m}^{-3}$ (based on the yearly averages of the 14 measurement locations). The yearly average gross α activity concentration in air dust is within the range of those from previous years, as illustrated in Figure 4.2. When comparing this value (yearly average of $3.7 \text{ Bq}\cdot\text{m}^{-3}$) with data collected before 2002, it should be noted that the Berthold measurements are 20% higher than the FAG measurements. The yearly average of the artificial β activity concentration does not deviate significantly from zero.


Since 2004, the analysis of the ambient dose equivalent rate has been based on a set of 153 stations. The yearly average ambient dose equivalent rate in 2016 was calculated using all 153 stations.

In 2016, the yearly average for the ambient dose equivalent rate was 81.1 nSv h^{-1} . This value, similar to the value of 2015, is significantly higher than the values measured before 2014, as can be seen in Figure 4.4. This increase of the ambient dose equivalent rate coincides with, and is attributable to, the replacement of the old Bitt RS03 proportional counter monitors with the new Saphymo XL 2-3 Geiger-Müller monitors.




Dots represent the locations of the aerosol monitors.

Figure 4.1 An impression of spatial variation in the average gross alpha activity concentration of (mainly) short-lived radionuclides in air dust




During the second half of 2002, the FAG monitors were replaced by Berthold monitors.

Figure 4.2 Yearly average gross alpha activity concentration of (mainly) short-lived radionuclides in air dust.



Dots represent the locations of the dose equivalent rate monitors.

Figure 4.3 An impression of spatial variation in the average ambient dose equivalent rate



During the course of 2015, most of the proportional counter monitors were replaced by Geiger-Müller monitors.

Figure 4.4 The yearly average ambient dose equivalent rate

Figure 4.5 shows the cosmogenic contribution to the effective dose rate, which is related to the ambient dose equivalent rate. Figure 4.5 also shows the influence of the 11-year solar cycle on the cosmogenic contribution.




Figure derived from data supplied by the Federal Aviation Administration [41].
Figure 4.5 Cosmogenic contribution to the effective dose rate (at sea level), influenced by the solar cycle: location 51° 26' north and 3° 43' east (in the south-western part of the Netherlands), air pressure 1019 hPa

5 Surface water and seawater

5.1 Introduction

The Rijkswaterstaat (RWS) regularly monitors the concentration of a large number of radionuclides in surface water and seawater. A representative part of the RWS's monitoring programme is presented here. A more detailed description of the monitoring programme, its underlying strategy and the results of radioactivity measurements in Dutch waters are reported elsewhere [42, 43, 44, 45].

The general monitoring strategy used for surface water is to monitor the inland and border-crossing water bodies of the Netherlands. Therefore, the locations mentioned in Table 5.1 are used for monitoring as they represent the major inland, incoming and outgoing waters of the Netherlands. The locations for seawater presented in this report have been chosen to represent the major areas of seawater.

The monitoring programme used for determining radionuclides in surface water and seawater is shown in Tables 5.1 and 5.2 and in Figure 5.1. Radionuclides were measured in water and in suspended solids. The samples were collected at equidistant times.

Since 2010, measurements in sediment have been added to the RWS monitoring programme, but the results are not presented in this report. These results are presented elsewhere [45]. The radionuclides were measured according to standard procedures [46, 47].

Table 5.1 Monitoring programme for the determination of radionuclides in surface water

Location	Parameter	Matrix	Monitoring frequency (per year)
IJsselmeer (Vrouwezand)	Gross α	Water	12
	Residual β	Water	12
	^3H	Water	7
	^{60}Co	Suspended solids	13
	^{131}I	Suspended solids	13
	^{137}Cs	Suspended solids	13
Noordzeekanaal (IJmuiden)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	13
	^{60}Co	Suspended solids	7
	^{131}I	Suspended solids	7
	^{137}Cs	Suspended solids	7

Continued on next page

Table 5.1 Continued

Location	Parameter	Matrix	Monitoring frequency (per year)
Nieuwe Waterweg (Maassluis)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	6
	^{90}Sr	Water	6
	^{226}Ra	Water	6
	^{60}Co	Suspended solids	13
	^{131}I	Suspended solids	13
	^{137}Cs	Suspended solids	13
	^{210}Pb	Suspended solids	6
Rhine (Lobith)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	13
	^{90}Sr	Water	7
	^{226}Ra	Water	7
	^{60}Co	Suspended solids	25
	^{131}I	Suspended solids	25
	^{137}Cs	Suspended solids	25
	^{210}Pb	Suspended solids	7
Scheldt (Schaar van Ouden Doel)	Gross α	Water	12
	Residual β	Water	12
	^3H	Water	7
	^{226}Ra	Water	7
	^{60}Co	Suspended solids	13
	^{131}I	Suspended solids	13
	^{137}Cs	Suspended solids	13
		^{210}Pb	Suspended solids
Meuse (Eijsden)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	13
	^{90}Sr	Water	7
	^{226}Ra	Water	7
	^{60}Co	Suspended solids	51
	^{131}I	Suspended solids	51
	^{137}Cs	Suspended solids	51
	^{210}Pb	Suspended solids	7
Kanaal Gent Terneuzen (Sas van Gent)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	6
	^{60}Co	Suspended solids	4
	^{131}I	Suspended solids	4
	^{137}Cs	Suspended solids	4
Haringvliet (Haringvlietsluis)	Gross α	Water	13
	Residual β	Water	13
	^3H	Water	7
	^{60}Co	Suspended solids	9
	^{131}I	Suspended solids	9
	^{137}Cs	Suspended solids	9

Table 5.2 Monitoring programme for the determination of radionuclides in seawater

Area	Location	Parameter	Matrix	Monitoring frequency (per year)
Coastal Area (KZ)	Noordwijk 2 ⁽¹⁾	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4
Southern North Sea (ZN)	Noordwijk 70 ⁽¹⁾	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4
		⁹⁰ Sr	Water	4
Central North Sea (CN)	Terschelling 235 ⁽¹⁾	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4
		⁹⁰ Sr	Water	4
Delta Coastal Waters (VD)	Schouwen 10 ⁽¹⁾	Gross α	Water	12
		Residual β	Water	12
		³ H	Water	12
		⁹⁰ Sr	Water	4
Western Scheldt (WS)	Vlissingen Boei	Gross α	Water	13
		Residual β	Water	13
		³ H	Water	13
		⁹⁰ Sr	Water	13
		¹³⁷ Cs	Suspended solids	4
		²¹⁰ Pb	Suspended solids	4
Eems-Dollard (ED)	Huibergat Oost	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4
Wadden Sea West (WW)	Marsdiep Noord	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4
Wadden Sea East (WO)	Dantziggat	Gross α	Water	4
		Residual β	Water	4
		³ H	Water	4

⁽¹⁾ Number indicates distance from shore. For example, 'Noordwijk 2' means Noordwijk 2 km offshore.




Figure 5.1 Overview of monitoring locations for the entire monitoring programme for surface water and seawater, as given by the Rijkswaterstaat

5.2 The results for surface water

The results of measurements of radioactivity in surface water are presented in Tables A11 and A12 and in Figures 5.2 to 5.19.

Gross α and residual β are indicative parameters. In general, gross alpha and beta analysis is used as a screening method to determine the total radioactivity present in the form of alpha and beta radiation, without regard to the identity of specific radionuclides.


The yearly average activity concentrations of gross α for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were 33, 109, 96, 49, 249, 34, 75 and 29 mBq·L⁻¹, respectively. The yearly average activity concentrations of residual β for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were 14, 21, 46, 29, 112, 24, 28 and 13 mBq·L⁻¹, respectively. The yearly average activity concentrations of gross α and residual β in 2016 were within the range of those in previous years.

Residual β in Noordzeekanaal, Nieuwe Waterweg and Scheldt has shown a change in trend since 1994, which was caused by a change in measuring technique that applies only to salt and brackish water [42]. This change in trend was therefore not seen for residual β in IJsselmeer, Rhine or Meuse.

The yearly average ³H activity concentrations for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were 2.4, 2.6, 3.2, 3.5, 10.8, 14.6, 1.6 and 5.0 Bq·L⁻¹, respectively. The yearly average ³H activity concentrations in 2016 were within the range of those in previous years. Elevated levels of ³H in the Rhine could have originated from several nuclear power plants or research reactors in Germany, France or Switzerland. Elevated levels of ³H in the Meuse could have originated from the nuclear power plants at Tihange (Belgium) or Chooz (France). Elevated levels of ³H in the Scheldt could have originated from the nuclear power plant at Doel (Belgium).

The nuclide ⁹⁰Sr is released into the environment by nuclear power plants and nuclear reprocessing plants. The yearly average ⁹⁰Sr activity concentrations for Nieuwe Waterweg, Rhine and Meuse were < 1.0, 3.1 and < 1.3 mBq·L⁻¹, respectively. The yearly average ⁹⁰Sr activity concentrations in 2016 were within the range of those in previous years.

The nuclide ²²⁶Ra is released into the environment by the ore-processing industry and transshipment. ²²⁶Ra in Nieuwe Waterweg and Scheldt might originate from these industries in the port areas of Rotterdam-Rijnmond and Antwerp, respectively. The yearly average ²²⁶Ra activity concentrations for Nieuwe Waterweg, Rhine, Scheldt and Meuse were 4.0, 8.6, 33.1 and 5.5 mBq·L⁻¹, respectively. The yearly average ²²⁶Ra activity concentrations in 2016 were within the range of those in previous years, with the exception of elevated ²²⁶Ra activity concentration in the Scheldt.



Average values are shown in cases of multiple measurements per month.
 Figure 5.2 Gross α activity concentrations for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet





Figure 5.3 Yearly average gross α activity concentrations



Average values are shown in cases of multiple measurements per month.
 Figure 5.4 Residual β activity concentrations for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet





Figure 5.5 Yearly average residual β activity concentrations



Average values are shown in cases of multiple measurements per month.
 Figure 5.6 ^3H activity concentrations for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet







Figure 5.7 Yearly average ^3H activity concentrations



Average values are shown in cases of multiple measurements per month.
 Figure 5.8 ⁹⁰Sr activity concentrations for Nieuwe Waterweg, Rhine and Meuse



Data are not available for Nieuwe Waterweg in 1995, 1996, 1999 and 2000.
 Figure 5.9 Yearly average ⁹⁰Sr activity concentrations



Average values are shown in cases of multiple measurements per month.
 Figure 5.10 ²²⁶Ra activity concentrations for Nieuwe Waterweg, Rhine, Scheldt and Meuse




Figure 5.11 Yearly average ²²⁶Ra activity concentrations

Nuclear power plants discharge radionuclides, including ^{60}Co and ^{137}Cs . ^{60}Co activity concentrations are higher in Meuse than elsewhere. ^{60}Co (and ^{137}Cs) in Meuse could have originated from the nuclear power plants at Tihange (Belgium) or Chooz (France).


The yearly average ^{60}Co activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were all $< 1 \text{ Bq}\cdot\text{kg}^{-1}$ except for Meuse ($10.4 \text{ Bq}\cdot\text{kg}^{-1}$).

The yearly average ^{137}Cs activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were 3.0, 6.2, 7.3, 10.3, 5.7, 10.1, 5.3 and $12.4 \text{ Bq}\cdot\text{kg}^{-1}$, respectively. In 2016, the yearly average ^{60}Co and ^{137}Cs activity concentrations in suspended solids were within the range of those in previous years.

The nuclide ^{131}I is released into the environment primarily by medical facilities. ^{131}I activity concentrations are higher in Noordzeekanaal and Meuse than elsewhere. ^{131}I in Noordzeekanaal and Meuse might originate from a sewage treatment plant in the port area of Westpoort and medical facilities in Belgium, respectively.

The yearly average ^{131}I activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet were < 1 , 36, < 2 , < 2 , < 1 , 6.2, 6.5 and $< 1.2 \text{ Bq}\cdot\text{kg}^{-1}$, respectively. In 2016, the yearly average ^{131}I activity concentrations in suspended solids were within the range of those in previous years.

The radionuclides ^{210}Po and ^{210}Pb originate from the uranium decay chain and are released by the ore-processing industry [42]. Since ^{210}Po is usually in equilibrium with ^{210}Pb in suspended solids, RWS reports only ^{210}Pb . ^{210}Pb in Nieuwe Waterweg and Scheldt might originate from these types of industries in the port areas of Rijnmond and Antwerp, respectively. The yearly average ^{210}Pb activity concentrations in suspended solids for Nieuwe Waterweg, Rhine, Scheldt and Meuse, were 97, 121, 80 and $140 \text{ Bq}\cdot\text{kg}^{-1}$, respectively. In 2016, the yearly average ^{210}Pb activity concentrations in suspended solids were within the range of those in previous years. For the Noordzeekanaal (at location IJmuiden), the presence of ^{131}I most likely originates from a sewage treatment facility. One of the contributions to the sewage waters is discharge from medical facilities. It has not been determined which changes over time might be related to the observed changes of activity concentrations over the years.



Average values are shown in cases of multiple measurements per month.
 Figure 5.12 ⁶⁰Co activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet





Figure 5.13 Yearly average ⁶⁰Co activity concentrations in suspended solids



Average values are shown in cases of multiple measurements per month.
 Figure 5.14 ¹³¹I activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet





Figure 5.15 Yearly average ¹³¹I activity concentrations in suspended solids



Average values are shown in cases of multiple measurements per month.
 Figure 5.16 ¹³⁷Cs activity concentrations in suspended solids for IJsselmeer, Noordzeekanaal, Nieuwe Waterweg, Rhine, Scheldt, Meuse, Kanaal Gent Terneuzen and Haringvliet





Figure 5.17 Yearly average ¹³⁷Cs activity concentrations in suspended solids



Average values are shown in cases of multiple measurements per month.
 Figure 5.18 ²¹⁰Pb activity concentrations in suspended solids




Figure 5.19 Yearly average ²¹⁰Pb activity concentrations in suspended solids

5.3 The results for seawater

The results of measurements of radioactivity in seawater are presented in Tables A13 and A14 and in Figures 5.20 to 5.31.

Gross α and residual β are indicative parameters [42]. In the first half of 2000, the background of the measuring equipment was unstable and higher than usual, which resulted in lower results. Therefore, yearly average concentrations of gross α in 2000 were based on data starting from the end of July 2000. Changes in the trend of gross α in the period 1985–1997 are explained elsewhere [42].

The yearly average activity concentrations of gross α in seawater for the Coastal Area, Southern North Sea, Central North Sea, Delta Coastal Waters, Western Scheldt, Eems-Dollard, Wadden Sea West and Wadden Sea East were 410, 350, 190, 340, 350, 210, 390 and 420 mBq·L⁻¹, respectively. The yearly average gross α activity concentrations in 2016 were within the range of those in previous years (Figure 5.21).

Residual β shows an apparent change in trend since 1994 (Figure 5.23). This was caused by a change in measuring technique that applies only to salt and brackish water [42]. The yearly average activity concentrations of residual β in seawater for the Coastal Area, Southern North Sea, Central North Sea, Delta Coastal Waters, Western Scheldt, Eems-Dollard, Wadden Sea West and Wadden Sea East were 59, 43, 34, 44, 110, 43, 68 and 140 mBq·L⁻¹, respectively. The yearly average residual β activity concentrations in 2016 were within the range of those in previous years (Figure 5.23).

Nuclear power plants discharge the radionuclides ³H and ¹³⁷Cs, among others. Nuclear fuel reprocessing plants discharge the radionuclides ³H and ⁹⁰Sr, among others. Discharges from the nuclear power plants at Doel (Belgium) and Borssele (Netherlands) are monitored in the Western Scheldt (WS) area. The impact of reprocessing plants at Sellafield (England) and Le Havre (France) is monitored in the Central North Sea (CN) and Southern North Sea (ZN) areas, respectively [42]. The impact of both sources (nuclear power and reprocessing plants) is monitored indirectly in the Delta Coastal Waters (VD) area.

The yearly average ³H activity concentrations in seawater for the Coastal Area, Southern North Sea, Central North Sea, Delta Coastal Waters, Western Scheldt, Eems-Dollard, Wadden Sea West and Wadden Sea East were 4.3, 2.6, 0.4, 4.7, 5.2, 3.8, 4.1 and 3.8 Bq·L⁻¹, respectively.

The yearly average ⁹⁰Sr activity concentrations in seawater for the Southern North Sea, Central North Sea, Delta Coastal Waters and Western Scheldt were 1.9, < 1.4, < 1.9 and <1.8 mBq·L⁻¹, respectively. The yearly average ³H and ⁹⁰Sr concentrations in 2016 were within the range of those in previous years (Figures 5.25 and 5.27).




Figure 5.20 Gross α activity concentrations in seawater for the Coastal Area (KZ), Southern North Sea (ZN), Central North Sea (CN), Delta Coastal Waters (VD), Western Scheldt (WS), Eems-Dollard (ED), Wadden Sea West (WW) and Wadden Sea East (WO)




Figure 5.21 Yearly average gross α activity concentrations




Figure 5.22 Residual β activity concentrations in seawater for the Coastal Area (KZ), Southern North Sea (ZN), Central North Sea (CN), Delta Coastal Waters (VD), Western Scheldt (WS), Eems-Dollard (ED), Wadden Sea West (WW) and Wadden Sea East (WO)




Figure 5.23 Yearly average residual β activity concentrations




Figure 5.24 ^3H activity concentrations in seawater for the Coastal Area (KZ), Southern North Sea (ZN), Central North Sea (CN), Delta Coastal Waters (VD), Western Scheldt (WS), Eems-Dollard (ED), Wadden Sea West (WW) and Wadden Sea East (WO)




Figure 5.25 Yearly average ^3H activity concentrations




Figure 5.26 ^{90}Sr activity concentrations in seawater for the Southern North Sea (ZN), Central North Sea (CN), Delta Coastal Waters (VD) and Western Scheldt (WS)




Figure 5.27 Yearly average ^{90}Sr activity concentrations

The radionuclides ^{210}Pb and ^{210}Po originate from the uranium decay chain and are released, for example, by the phosphate-processing industry and production platforms for oil and gas [42]. The phosphate-processing industry has not been operational in the Netherlands since 2012. Since ^{210}Po is usually in equilibrium with ^{210}Pb in suspended solids, RWS reports only ^{210}Pb (as in surface water). In cases in which a strong increase in the gross α value is noticed, however, ^{210}Po is determined as well. Discharges via the main rivers are monitored in the Coastal Area (KZ). Discharges from the ore- and phosphate-processing industries in Belgium and the Netherlands are monitored in the Western Scheldt (WS) area. Discharges from Delfzijl, Eemshaven and plants in Germany are monitored in the Eems-Dollard (ED) area. The impact of these discharges, together with activity originating from the North Sea, is monitored indirectly in the Wadden Sea (WW and WO) area.

Since 2009, Wadden Sea West has replaced Wadden Sea East as a monitoring location. Since 2014, the monitoring of suspended solids has been discontinued in the Coastal Area, Eems-Dollard and Wadden Sea West.

The yearly average ^{137}Cs and ^{210}Pb activity concentrations in suspended solids for Western Scheldt were 3.3 and $62 \text{ Bq}\cdot\text{kg}^{-1}$, respectively. The yearly average ^{137}Cs and ^{210}Pb activity concentrations in 2016 were within the range of those in previous years (Figures 5.29 and 5.31).





Figure 5.28 ^{137}Cs activity concentrations in suspended solids in seawater for the Western Scheldt (WS)



Since 2009, Wadden Sea West (WW) has replaced Wadden Sea East (WO) as a monitoring location. Since 2014, the monitoring of suspended solids has been discontinued in the Coastal Area (KZ), Eems-Dollard (ED) and Wadden Sea West (WW).

Figure 5.29 Yearly average ¹³⁷Cs activity concentrations in suspended solids





Figure 5.30 ²¹⁰Pb activity concentrations in suspended solids in seawater for the Western Scheldt (WS)



Since 2009, Wadden Sea West (WW) has replaced Wadden Sea East (WO) as a monitoring location. Since 2014, the monitoring of suspended solids has been discontinued in the Coastal Area (KZ), Eems-Dollard (ED) and Wadden Sea West (WW).

Figure 5.31 Yearly average ²¹⁰Pb activity concentrations in suspended solids

6 Water for human consumption

6.1 Introduction

In addition to the Recommendation on the Application of Article 36 of the Euratom Treaty [1], requirements for the monitoring of drinking water are given in Council Directive 2013/51/EC [48]. According to this directive, the parameters ^3H and the total indicative dose should be monitored. Screening methods for gross α and gross β activity concentrations may be used to monitor the total indicative dose. If the gross α and gross β activity concentrations are less than the screening levels of 0.1 and 1.0 $\text{Bq}\cdot\text{L}^{-1}$, respectively, it can be assumed that the total indicative dose is less than the parametric value of 0.1 $\text{mSv}\cdot\text{year}^{-1}$ [49, 50, 51].

In the Netherlands, drinking water production stations monitor untreated water and treated water for ^3H , gross α , gross β and residual β activity concentrations. The monitoring frequency per location ranges from 1 to 26 times per year, depending on the volume of water produced. The activity concentrations are averaged over a year for each production station.

6.2 Results

The results for 2016 are presented in Table 6.1. For gross α , ^3H , gross β and residual β , several hundred analyses were performed at a number of production stations; the number of production stations varied between 168 and 186.

Table 6.1 Drinking water analyses in 2016

Parameter	Gross α	^3H	Residual β	Gross β
Average value ⁽¹⁾	< 0.06 $\text{Bq}\cdot\text{L}^{-1}$	< 4.0 $\text{Bq}\cdot\text{L}^{-1}$	< 0.1 $\text{Bq}\cdot\text{L}^{-1}$	< 0.1 $\text{Bq}\cdot\text{L}^{-1}$
No. of all production stations	183	180	168	186
No. of all analyses	388	448	390	428
Maximum value ⁽²⁾	0.44 $\text{Bq}\cdot\text{L}^{-1}$	16.5 $\text{Bq}\cdot\text{L}^{-1}$	< 0.2 $\text{Bq}\cdot\text{L}^{-1}$	0.6 $\text{Bq}\cdot\text{L}^{-1}$
No. of production stations ⁽³⁾	1	1	11	1
No. of analyses ⁽⁴⁾	1	13	103	1

⁽¹⁾ Activity concentration averaged over all production stations.

⁽²⁾ Maximum value of activity concentration averaged per production station.

⁽³⁾ Number of production stations with maximum value.

⁽⁴⁾ Number of analyses performed per production station that led to maximum value.

In 2016, the gross α activity concentration in untreated water used for drinking water production, averaged per production station, exceeded the screening level of 0.1 $\text{Bq}\cdot\text{L}^{-1}$ at 6 of the 183 production stations (in 19 of the 388 analyses). The measured radioactivity levels do not, however, pose a threat to public health. Further investigation into these slightly elevated levels in untreated water revealed that the gross α activity concentration in associated finished drinking water was well below the screening level.

For ^3H , gross β and residual β , the results were within the range of those in previous years [7, 20, 52, 53, 54, 55, 56, 57, 58, 59, 20, 60, 61, 62,

63]. Since ^{40}K was almost absent, for most stations there was no significant difference between average gross β and residual β activity concentrations. The gross β activity concentrations were below the screening level of $1.0 \text{ Bq}\cdot\text{L}^{-1}$ and the ^3H activity concentrations were below the parametric value of $100 \text{ Bq}\cdot\text{L}^{-1}$ [48, 50, 51].

Following Council Directive 2013/51/EC [48] it is not necessary to incorporate ^{222}Rn in the yearly Dutch drinking water monitoring programme if it is clearly demonstrated that all data are well below accepted levels. The routine determination of gross α , gross β and tritium covers most radioactivity parameters and gives an accurate estimation of the dose (indicative dose).

In 2015, a survey was carried out to determine radon activity in Dutch water [64]. The results of the 2015 survey have been summarised in [63]: the parametric value of $100 \text{ Bq}\cdot\text{L}^{-1}$ of ^{222}Rn was not exceeded. The highest concentration of ^{222}Rn in groundwater was $16.7 \text{ Bq}\cdot\text{L}^{-1}$, with consistently lower amounts of ^{222}Rn in finished drinking water than in groundwater. All observed ^{222}Rn activities originated from unsupported ^{222}Rn , i.e. without any ^{226}Ra present in the water phase.

7 Milk

7.1 Introduction

RIKILT Wageningen UR monitors radioactivity in milk on a weekly basis, mainly via the National Monitoring Network of Radioactivity in Food (Landelijk Meetnet Radioactiviteit in Voedsel, LMRV). The LMRV has been set up as an emergency network for monitoring relatively high contamination levels in case of an accident. The LMRV consists of 48 low-resolution γ -spectrometers (NaI-detectors) located throughout the Netherlands, 22 of which are located at dairy factories.

7.2 Results

The results of the weekly samples of cow's milk taken from all locations are combined into a monthly average for the whole country. The monthly averages for 2016 are presented in Table 7.1. Figure 7.1 shows the spatial variation of the yearly average ^{40}K concentrations per region and the distribution of the sampling locations across the Netherlands.

Table 7.1 Monthly average activity concentrations in cow's milk in 2016.

Month	Number of samples	^{40}K ⁽¹⁾ Bq·kg ⁻¹	^{60}Co ⁽²⁾ Bq·kg ⁻¹	^{131}I ⁽²⁾ Bq·kg ⁻¹	^{134}Cs ⁽²⁾ Bq·kg ⁻¹	^{137}Cs ⁽²⁾ Bq·kg ⁻¹
January	45	51.9 ± 10.0	< 1.4	< 0.6	< 0.6	< 0.5
February	51	50.6 ± 13.9	< 1.4	< 0.6	< 0.6	< 0.5
March	51	50.7 ± 11.0	< 1.4	< 0.6	< 0.6	< 0.5
April	41	51.1 ± 12.3	< 1.4	< 0.6	< 0.6	< 0.5
May	55	50.9 ± 13.4	< 1.4	< 0.6	< 0.6	< 0.5
June	55	46.9 ± 9.1	< 1.4	< 0.6	< 0.6	< 0.5
July	45	52.1 ± 14.1	< 1.4	< 0.6	< 0.6	< 0.5
August	49	50.1 ± 13.1	< 1.4	< 0.6	< 0.6	< 0.5
September	43	53.1 ± 12.3	< 1.4	< 0.6	< 0.6	< 0.5
October	39	54.1 ± 14.7	< 1.4	< 0.6	< 0.6	< 0.5
November	48	51.3 ± 12.2	< 1.4	< 0.6	< 0.6	< 0.5
December	59	52.7 ± 11.3	< 1.4	< 0.6	< 0.6	< 0.5
Average	581 ⁽³⁾	51.2 ± 12.3	< 1.4	< 0.6	< 0.6	< 0.5

(1) Uncertainty is given as 1σ .

(2) Calculated minimal detectable activity concentrations for the respective radionuclides, based on 1 litre Marinelli beaker measurements on the Food Monitor Systems.

(3) Yearly total.


In none of the samples were anthropogenic γ -emitters were measured above the minimal detectable activity, as is shown in Table 7.1, so the limit of 370 Bq·kg⁻¹ for the radiocesium activity (sum of ^{134}Cs and ^{137}Cs) set by the European Union [65, 66] was not exceeded. The activity concentration of the natural radionuclide ^{40}K is given as a reference value. The yearly average concentration was 51.2 ± 12.3 Bq·kg⁻¹. This value is within the range of those found in previous years.

Additionally, 11 goat's milk samples were analysed. As in cow's milk, anthropogenic γ -emitters were not measured above the minimal detectable activity. The yearly average ^{40}K concentration in these samples was 50.1 ± 7.6 Bq·kg⁻¹. This value is lower than those found in previous years.

In addition to the LMRV samples, 47 milk samples (45 cow's milk and 2 goat's milk samples) were analysed for a range of γ -emitters on a high-resolution gamma spectrometer in the RIKILT laboratory in Wageningen. The samples were collected across the Netherlands. None of the samples showed any anthropogenic gamma activity above the minimal detectable activity ($<1 \text{ Bq}\cdot\text{kg}^{-1}$ for ^{137}Cs in 0.5 L Marinelli beakers). The average concentration found for the natural radionuclide ^{40}K in the 45 cow's milk samples was $45.4 \pm 9.7 \text{ Bq}\cdot\text{kg}^{-1}$; for the 2 goat's milk samples the average was $66.9 \pm 7.4 \text{ Bq}\cdot\text{kg}^{-1}$.

The same 47 raw milk samples were analysed for the presence of the β -emitter ^{90}Sr using low-level liquid scintillation counting (LSC). The ^{90}Sr activity concentration was below the minimal detectable activity ($0.2 \text{ Bq}\cdot\text{kg}^{-1}$) in all samples taken, so none of the samples exceeded the set limit of $125 \text{ Bq}\cdot\text{kg}^{-1}$ used in new emergency exposure situations [67]. No limit for ^{90}Sr has been set for existing exposure situations as defined in [68].

RIKILT also monitors raw milk specifically for export certification. For this, samples were analysed for ^{137}Cs and ^{90}Sr . All results were below minimum detectable activities as well.



Based on data provided by dairy factories.

Figure 7.1 Impression of the spatial variation of ^{40}K activity concentrations ($\text{Bq}\cdot\text{kg}^{-1}$) in cow's milk

8 Food

8.1 Introduction

The Netherlands Food and Consumer Product Safety Authority (NVWA) performs measurements on finished products from retail shops, wholesale produce auctions and distribution centres, while RIKILT Wageningen UR performs measurements on samples from earlier stages in the food production chain.

The measurements performed on food by the NVWA were carried out according to standard procedures [69, 70]. Since 2005, the NVWA has monitored activity concentrations in a 'mixed diet' every year by sampling and measuring separate ingredients. Over a period of four weeks in 2016, 372 samples were taken from retail shops, wholesale produce auctions and distribution centres, including 49 samples of honey [71]. Though honey is not considered to be part of the mixed diet, samples are taken each year because it regularly contains higher levels of radioactivity (mainly ^{137}Cs).

The separate ingredients were categorised into the following product groups: grain and grain products, vegetables and mushrooms, fruit and fruit products, milk and dairy products, meat and meat products, game and poultry, salads, oil and butter, honey, tea, mineral water, and fish. The 2016 results are presented in Table 8.1. All three samples containing ^{137}Cs above the minimum detectable activity were dried mushrooms, with the highest measured value of 1,019 Bq/kg of ^{137}Cs . As mushrooms are not consumed in their dry form, the activity concentration in its consumable form is compared to the set limit of 600 Bq/kg, according to EC regulation No. 73/2008 [66]. This is achieved by multiplying by a reduction factor that can vary between 0.1 and 0.25. None of the values exceeded the set limit of 600 Bq·kg⁻¹ (or 370 Bq·kg⁻¹ for milk and dairy products) [65, 66] so no further action was taken.

In 2016, RIKILT Wageningen UR analysed radioactivity in food products as part of the governmental monitoring programme. Samples were taken throughout the year and measurements were carried out according to standard procedures. A total of 1,702 food samples were analysed for the presence of γ -emitters. The results are presented in Table 8.2. None of the samples exceeded the set limit for radiocesium activity (sum of ^{134}Cs and ^{137}Cs) of 600 Bq·kg⁻¹ (for food) or 370 Bq·kg⁻¹ (for dairy products).

Of these food samples, 179 samples were additionally analysed for ^{90}Sr content. The results are presented in Table 8.3. These results are well below the set limit for new emergency exposure situations of 750 Bq·kg⁻¹ for major food products [67]. No limit for ^{90}Sr has been set for existing exposure situations as defined in [68].

RIKILT also monitors food specifically for export certification. For this, samples were analysed for ^{137}Cs and ^{90}Sr . All results were below the limits set for ^{137}Cs and below minimal detectable activity for ^{90}Sr .

8.2 Results for honey

In total, 49 samples of honey were analysed by the NVWA [71]. No sample of honey contained ^{137}Cs above the minimum detectable activity of $5 \text{ Bq}\cdot\text{kg}^{-1}$. Radiocesium activity was below the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$ [65, 66].

8.3 Results for vegetables and mushrooms

In the product group 'vegetables and mushrooms' analysed by the NVWA, concentrations above the minimum detectable activity of $5 \text{ Bq}\cdot\text{kg}^{-1}$ of ^{137}Cs were detected in three samples of dried mushrooms: 58, 222 and $1,019 \text{ Bq}\cdot\text{kg}^{-1}$ of ^{137}Cs . As mushrooms are not consumed in their dry form, the activity concentration in its consumable form is compared to the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$. None of the values exceeded the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$ [65, 66].

8.4 Results for game and poultry

In the product group 'game and poultry' analysed by RIKILT Wageningen UR, 24 samples of game contained ^{137}Cs . The activity varied from 5 up to $43 \text{ Bq}\cdot\text{kg}^{-1}$, all below the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$ [65, 66].

8.5 Results for average daily intake

The measured concentrations of ^{90}Sr , ^{134}Cs and ^{137}Cs in food in $\text{Bq}\cdot\text{kg}^{-1}$ were converted to an average daily intake value per person per day ($\text{Bq}\cdot\text{day}^{-1}$) using food consumption patterns, according to the method described in Appendix B. From these intake values, a contribution to the effective yearly dose was calculated using standard dose conversion coefficients for ingestion.

The average daily intake per person of ^{134}Cs , ^{137}Cs and ^{90}Sr is estimated at < 5 , < 7 , $< 5 \text{ Bq}\cdot\text{day}^{-1}$, respectively. These estimates are mainly based on the minimal detectable activities for these radionuclides in the different food categories, as shown in Tables 8.1 to 8.3.

The contribution to the effective yearly dose calculated from these average daily intake values is $< 0.1 \text{ mSv}$. The actual daily intake (and following dose contribution) is probably much lower.

Table 8.1 Results of 2016 analysis of food for ^{134}Cs and ^{137}Cs as measured by the Netherlands Food and Consumer Product Safety Authority

Product	Number of samples	^{134}Cs ⁽¹⁾ Bq·kg ⁻¹	^{137}Cs ⁽¹⁾ Bq·kg ⁻¹
Grain and grain products	31	< 5 (0)	< 5 (0)
Vegetables and mushrooms	84	< 5 (0)	58, 222, 1019 (3)
Fruit and fruit products	10	< 5 (0)	< 5 (0)
Milk and dairy products	50	< 5 (0)	< 5 (0)
Meat and meat products	32	< 5 (0)	< 5 (0)
Game and poultry	26	< 5 (0)	< 5 (0)
Salads	17	< 5 (0)	< 5 (0)
Oil and butter	30	< 5 (0)	< 5 (0)
Honey	49	< 5 (0)	< 5 (0)
Tea	3	< 5 (0)	< 5 (0)
Mineral water	18	< 5 (0)	< 5 (0)
Fish	22	< 5 (0)	< 5 (0)

⁽¹⁾ Number of samples above the given reporting limit is shown in brackets.

Table 8.2 Results of 2016 analysis of food for ^{134}Cs and ^{137}Cs as measured by RIKILT Wageningen UR

Product	Number of samples	^{134}Cs ⁽¹⁾ Bq·kg ⁻¹	^{137}Cs ⁽¹⁾ Bq·kg ⁻¹
Vegetables and fruits	287	< 5 (0)	< 5 (0)
Meat and meat products	568	< 5 (0)	< 5 (0)
Game and poultry	398	< 5 (0)	5–43 (24) ⁽²⁾
Eggs	156	< 5 (0)	< 5 (0)
Fish and seafood products	201	< 5 (0)	< 5 (0)
Ready meals	45	< 5 (0)	< 5 (0)

⁽¹⁾ Number of samples above the given detection limit is shown in brackets.

⁽²⁾ Range of activity concentrations above the minimum detectable activity level.

Table 8.3 Results of 2016 analysis of food for ^{90}Sr as measured by RIKILT Wageningen UR

Product	Number of samples	^{90}Sr ⁽¹⁾ Bq·kg ⁻¹
Vegetables and fruits	23	< 5 (0)
Meat and meat products	15	< 5 (0)
Game and poultry	19	< 5 (0)
Eggs	6	< 5 (0)
Fish and seafood products	24	< 5 (0)
Ready meals	45	< 5 (0)

⁽¹⁾ Number of samples above the minimum detectable activity level is shown in brackets.

9 Grass & feed

The National Monitoring Network of Radioactivity in Food (Landelijk Meetnet Radioactiviteit in Voedsel, LMRV), referred to in Chapter 7, is an important monitoring network used in cases of a nuclear or radiological emergency, as described in the National Crisis Management Plan for Radiation Incidents (Nationaal Crisisplan Stralingsincidenten, NCS). In addition to measuring radioactivity levels in milk and food samples, the network is used to measure radioactivity levels in grass samples. For this purpose, reference pastures and fields have been designated across the Netherlands in proximity to the companies and organisations that participate in the LMRV. In this way, the extent of radioactive deposition can be assessed rapidly by the LMRV in the event of a nuclear or radiological incident.

It is important to have accurate and recent information on the natural background levels of radioactivity in grass to compare with samples analysed during a nuclear or radiological incident. For this reason, all LMRV locations are requested to take a grass sample every year from their reference pasture or field according to a standardised protocol, and to measure this sample using the food monitoring system.

In 2016, 36 grass samples were taken at 19 locations and measured on the food monitoring system. None of the grass samples taken contained artificial radionuclides above the minimal detectable activities. The minimal detectable activities were approximately $20 \text{ Bq}\cdot\text{m}^{-2}$ (i.e. with a yield of 250 gram grass per m^2) for the artificial radionuclides ^{60}Co , ^{131}I , ^{132}Te , ^{134}Cs and ^{137}Cs . Natural ^{40}K was found in all samples. In some samples, natural radionuclides from the uranium and thorium decay chains deposited during rainfall were measured as well.

The results of the grass measurements can be plotted on a map of the Netherlands, resulting in a deposition map that can be used in the event of contamination in an emergency situation. The spatial variation of the natural ^{40}K activity in grass is shown in Figure 9.1. This variation in ^{40}K activity can be attributed to factors such as fertilisation, grass species, length of stalk and soil type.

In addition, 462 feed samples were analysed for γ -emitters as part of the monitoring programme of RIKILT Wageningen UR. No artificial radioactivity was found in these samples; the results for ^{134}Cs and ^{137}Cs were all lower than the minimal detectable activity ($2 \text{ Bq}\cdot\text{kg}^{-1}$).




Figure 9.1 Impression of spatial variation of ^{40}K activity in grass in $\text{Bq}\cdot\text{m}^{-2}$, as measured in 2016

10 The Borssele nuclear power plant

The Nuclear Research & Consultancy Group (NRG) is commissioned by Elektriciteits-Produktie­maatschappij Zuid-Nederland (N.V. EPZ) to perform monthly measurements on environmental samples taken in the vicinity of the Borssele nuclear power plant (owned by N.V. EPZ). Samples are taken to monitor the compartments of air, water and soil [72]. A more detailed description of the monitoring programme and underlying strategy can be found elsewhere [73]. The 2016 monitoring programme for environmental samples is outlined in Table 10.1 and Figure 10.1. Radionuclides were determined in air dust, grass, soil, water, suspended solids, seaweed and sediment.

Table 10.1 Monitoring programme for environmental samples taken near the Borssele nuclear power plant in 2016

Matrix	Location	Parameter	Monitoring frequency (per year)
Air dust	21, 22, 23, 27 and 29	gross α , gross β γ -emitters ⁽¹⁾	12 (11 for location 29) 12 ⁽²⁾
Grass	21, 22, 23, 27 and 29	γ -emitters ⁽³⁾	12 ⁽²⁾
Soil	O1, O2, O3 and O4 ⁽⁴⁾	γ -emitters ⁽⁵⁾	1
Water	1, 2, 3 and 4	residual β , ³ H	12
Suspended solids	1, 2, 3 and 4	gross β	12
Seaweed	1, 2, 3 and 4	γ -emitters ⁽³⁾	12 ⁽²⁾
Sediment	1, 2, 3 and 4	γ -emitters ⁽³⁾	12 ⁽²⁾

The location numbers correspond to the location numbers given in Figure 10.1.


⁽¹⁾ γ -spectroscopic analysis of specific γ -emitting radionuclides: ⁶⁰Co, ¹³⁷Cs, naturally occurring radionuclides and elemental and organically bound ¹³¹I.

⁽²⁾ Analysis was performed on a combined sample of monthly samples taken from all four or five locations.

⁽³⁾ γ -spectroscopic analysis of specific γ -emitting radionuclides: ⁶⁰Co, ¹³¹I and ¹³⁷Cs.

⁽⁴⁾ The four locations where samples were taken near the outlet are not shown in Figure 10.1.

⁽⁵⁾ γ -spectroscopic analysis of specific γ -emitting radionuclides: ⁵⁴Mn, ⁶⁰Co, ¹³⁴Cs and ¹³⁷Cs.



The numbers given in Table 10.1 correspond with the locations on the map.

Figure 10.1 Overview of monitoring locations for the monitoring programme conducted by NRG near the Borssele nuclear power plant

10.2 Air

The results for gross α and β activity concentrations in air dust are presented in Tables A15 and A16. Due to large uncertainties caused by variations in the amount of dust on the filters, gross α activity concentrations in air dust should be regarded as indicative values [5]. The period between sampling and analysis was at least 5 days, which is long relative to the decay time of the short-lived decay products of ^{222}Rn and ^{220}Rn . This is to ensure that these naturally occurring decay products do not contribute to the measured α and β activity concentrations.

The 2016 yearly averages of the gross α and β activity concentrations of long-lived radionuclides were within the range of the results from previous years, as illustrated in Figures 10.2 and 10.3.

The results for the radionuclides considered in the γ -spectroscopic analysis are given in Table A17.




Figure 10.2 Yearly average gross α activity concentrations in air dust at five locations near the Borssele nuclear power plant (see Figure 10.1)




Figure 10.3: Yearly average gross β activity concentrations in air dust at five locations near the Borssele nuclear power plant (see Figure 10.1)

10.3 Soil and grass

The results for the radionuclides considered in the γ -spectroscopic analysis of grass and soil are given in Tables A18 and A19. The four soil samples were taken near the outlet of the nuclear power plant. In 2016,

the yearly average concentrations (in grass) and the yearly concentrations (in soil) of ^{54}Mn , ^{60}Co , ^{134}Cs and ^{137}Cs were within the range of those in previous years [20, 56, 57, 58, 59, 60, 61, 62, 63]. Except for ^{137}Cs in soil all results were below the minimum detectable activities. The yearly concentrations of ^{137}Cs in soil are shown in Figure 10.4.




Figure 10.4 ^{137}Cs activity concentrations of yearly soil sample at four locations near the outlet of the Borssele nuclear power plant (see Figure 10.1)

10.4 Water

The results for residual β and ^3H activity concentrations in surface water and gross β activity concentrations in suspended solids in the Western Scheldt area are presented in Tables A20, A21 and A22.

In 2016, the yearly averages of the residual β concentrations in surface water were within the range of the results from previous years, as illustrated in Figure 10.5. Since 2012, the ^3H activity concentrations in water have been significantly lower than those in previous years, as illustrated in Figure 10.6. Since 2012, the gross β activity concentrations in suspended solids have been somewhat higher than those in previous years, as illustrated in Figure 10.7. The changes in trends of ^3H and gross β activity concentrations were investigated. For gross β there has been a change in the counting efficiency used, which is a plausible explanation for the change in trend of the gross β activity concentration. For ^3H no significant changes in analysis procedures have been found.




Figure 10.5 Yearly average residual β activity concentrations in surface water in the Western Scheldt area at four locations near the Borssele nuclear power plant (see Figure 10.1)




Figure 10.6 Yearly average ^3H activity concentrations in surface water in the Western Scheldt area at four locations near the Borssele nuclear power plant (see Figure 10.1)




Figure 10.7 Yearly average gross β activity concentrations in suspended solids in the Western Scheldt area at four locations near the Borssele nuclear power plant (see Figure 10.1)




Figure 10.8 Yearly average ¹³⁷Cs activity concentrations in sediment based on combined samples at four locations near the Borssele nuclear power plant.

The results for the radionuclides considered in the γ -spectroscopic analysis of seaweed and sediment are given in Tables A23 and A24. Except for ¹³⁷Cs in sediment, all results were below the minimum detectable activities. The yearly concentrations of ¹³⁷Cs in sediment are shown in Figure 10.8.

11 Conclusions

In 2016, the Netherlands were in compliance with the Euratom recommendations on annual measurement of radioactivity in the environment and in food.

The gross α activity concentration in untreated water used for drinking water production, averaged per production station, exceeded the screening level ($0.1 \text{ Bq}\cdot\text{L}^{-1}$) at 6 of the 183 production stations (in 19 of the 388 analyses). The measured radioactivity levels do not, however, pose a threat to public health. Further investigation into these slightly elevated levels in untreated water revealed that the gross α activity concentration in associated finished drinking water was well below the screening level.

Radioactivity was measured in over 2,000 food products, 27 samples of which contained ^{137}Cs . Three samples of dried mushrooms contained ^{137}Cs (58, 222 and $1,109 \text{ Bq}\cdot\text{kg}^{-1}$). As mushrooms are not consumed in their dry form, the activity concentration in its consumable form is compared to the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$. Twenty-four samples of game and poultry contained ^{137}Cs and the activity varied from 5 to $43 \text{ Bq}\cdot\text{kg}^{-1}$. Radioactivity levels in none of the samples were above the set limit of $600 \text{ Bq}\cdot\text{kg}^{-1}$ (or $370 \text{ Bq}\cdot\text{kg}^{-1}$ for milk and dairy products) for the activity of radiocesium (sum of ^{134}Cs and ^{137}Cs).

The measured concentrations of ^{90}Sr , ^{134}Cs and ^{137}Cs in food in $\text{Bq}\cdot\text{kg}^{-1}$ were converted to an average daily intake value per person per day ($\text{Bq}\cdot\text{day}^{-1}$) using food consumption patterns. The average daily intake per person of ^{134}Cs , ^{137}Cs and ^{90}Sr is < 5 , < 7 , $< 5 \text{ Bq}\cdot\text{day}^{-1}$, respectively. The contribution to the effective yearly dose calculated from these values is $< 0.1 \text{ mSv}$. The actual daily intake (and following dose contribution) is probably much lower.

All other radioactivity measurements were within the range of those in previous years.

Appendix A Tables of results

Table A1 Weekly average gross α and gross β activity concentrations in air dust sampled with the Snow White high-volume sampler at RIVM in 2016

Week ⁽¹⁾ number	Gross α ⁽²⁾ mBq.m ⁻³	Gross β mBq.m ⁻³	Week ⁽¹⁾ number	Gross α ⁽²⁾ mBq.m ⁻³	Gross β mBq.m ⁻³
1	0.020	0.40 ± 0.04	27	0.011	0.167 ± 0.018
2	0.008	0.159 ± 0.017	28	0.017	0.27 ± 0.03
3	0.017	0.27 ± 0.03	29	0.022	0.38 ± 0.04
4	0.021	0.30 ± 0.03	30	0.029	0.52 ± 0.05
5	0.009	0.131 ± 0.014	31	0.011	0.144 ± 0.015
6	0.010	0.20 ± 0.02	32	0.015	0.25 ± 0.03
7	0.018	0.34 ± 0.04	33	0.012	0.23 ± 0.02
8	0.015	0.20 ± 0.02	34	0.025	0.39 ± 0.04
9	0.014	0.21 ± 0.02	35	0.023	0.48 ± 0.05
10	0.014	0.19 ± 0.02	36	0.014	0.23 ± 0.02
11	0.020	0.44 ± 0.05	37	0.040	0.97 ± 0.10
12	0.016	0.27 ± 0.03	38	0.031	0.79 ± 0.08
13	0.023	0.21 ± 0.02	39	0.022	0.61 ± 0.06
14	0.029	0.38 ± 0.04	40	0.011	0.24 ± 0.03
15	0.015	0.24 ± 0.03	41	0.011	0.28 ± 0.03
16	0.017	0.20 ± 0.02	42	0.038	0.77 ± 0.08
17	0.010	0.161 ± 0.017	43	0.030	0.60 ± 0.06
18 ⁽³⁾	0.012	0.23 ± 0.02	44	0.023	0.42 ± 0.04
19 ⁽³⁾	0.0075	0.50 ± 0.05	45	0.023	0.33 ± 0.03
20	0.010	0.42 ± 0.04	46	0.028	0.46 ± 0.05
21	0.010	0.21 ± 0.02	47	0.011	0.25 ± 0.03
22	0.010	0.47 ± 0.05	48 ⁽³⁾	0.015	0.24 ± 0.03
23	0.014	0.45 ± 0.05	49 ⁽³⁾	0.034	0.65 ± 0.05
24	0.029	0.42 ± 0.04	50	0.038	0.80 ± 0.08
25	0.018	0.23 ± 0.02	51	0.032	0.70 ± 0.07
26	0.014	0.25 ± 0.03	52	0.017	0.174 ± 0.018
Average				0.019	0.362 ± 0.006 ⁽⁴⁾
SD ⁽⁵⁾				0.009	0.19

⁽¹⁾ The precise sampling period is given in Table A3.

⁽²⁾ Values are indicative due to large uncertainties caused by variations in the amount of dust on the filters [5].

⁽³⁾ The sampling period deviated from the regular 7-day sampling period.

⁽⁴⁾ The uncertainty in the yearly average is equal to the square root of the sum of the squared weekly uncertainties divided by the number of weeks. Uncertainties are given as 1σ .

⁽⁵⁾ SD is the standard deviation of the weekly results.

Table A2 Detection limits ($\mu\text{Bq}/\text{m}^3$) in the residue measurement of air dust sampled during a 7-day sampling period with the Snow White high-volume sampler at RIVM in 2016

Nuclide	Detection limit $\mu\text{Bq}\cdot\text{m}^{-3}$
^7Be	2.0
^{22}Na	0.2
^{60}Co	0.1
^{131}I	1.7 ⁽¹⁾
^{137}Cs	0.1
^{210}Pb	3.7

Measurements were carried out on ash residue using a well-type detector, with a 7-day delay between sampling and the start of measurement. The sample volume was about 125,000 m³. Between January 2000 and July 2009, the detection limits were higher than they had been before 2000 [74] due to a different detector set-up. The detector set-up was changed again in the second half of 2009, including a change in counting time from 100,000 seconds to 178,200 seconds. Detection limits were therefore lower from July 2009 until January 2011, when a change in the high-volume sampler (and consequently the sample volume) resulted in a further reduction of the detection limits.

⁽¹⁾ The detection limit is given for the filter measurement on the coaxial detector (4-day delay time, 100,000 seconds counting time). Due to the sample preparation procedure, the volatile radionuclide ^{131}I cannot be determined in the residue measurement on the well-type detector.

Table A3 Weekly average ^7Be , ^{137}Cs and ^{210}Pb activity concentrations in air dust sampled with the Snow White high-volume sampler at RIVM in 2016

Week number	Period	^7Be $\mu\text{Bq}\cdot\text{m}^{-3}$	^{137}Cs $\mu\text{Bq}\cdot\text{m}^{-3}$	^{210}Pb $\mu\text{Bq}\cdot\text{m}^{-3}$
1	31/12–07/01	2,030 ± 150	0.35 ± 0.04	450 ± 30
2	07/01–14/01	2,300 ± 170	0.08 ± 0.02	117 ± 9
3	14/01–21/01	2,520 ± 180	0.27 ± 0.03	226 ± 17
4	21/01–28/01	2,130 ± 160	0.17 ± 0.03	280 ± 20
5	28/01–04/02	2,210 ± 160	< 0.09	81 ± 6
6	04/02–11/02	2,250 ± 160	0.04 ± 0.02	130 ± 10
7	11/02–18/02	1,930 ± 140	0.35 ± 0.04	320 ± 20
8	18/02–25/02	1,510 ± 110	0.06 ± 0.02	131 ± 10
9	25/02–03/03	1,980 ± 150	0.19 ± 0.03	128 ± 9
10	03/03–10/03	1,940 ± 140	0.19 ± 0.03	140 ± 10
11	10/03–17/03	2,430 ± 180	0.53 ± 0.05	470 ± 40
12	17/03–24/03	2,800 ± 200	0.36 ± 0.04	248 ± 18
13	24/03–31/03	3,100 ± 200	0.079 ± 0.018	122 ± 9
14	31/03–07/04	3,800 ± 300	0.16 ± 0.03	290 ± 20
15	07/04–14/04	3,100 ± 200	0.13 ± 0.02	181 ± 13
16	14/04–21/04	2,370 ± 170	0.08 ± 0.03	128 ± 9
17	21/04–28/04	2,540 ± 190	0.13 ± 0.03	116 ± 9
18 ⁽¹⁾	28/04–04/05	3,200 ± 200	0.08 ± 0.02	157 ± 12
19 ⁽¹⁾	04/05–12/05	5,000 ± 400	0.67 ± 0.06	530 ± 40
20	12/05–19/05	5,600 ± 400	0.47 ± 0.04	330 ± 20
21	19/05–26/05	2,480 ± 180	0.11 ± 0.03	175 ± 13
22	26/05–02/06	3,300 ± 200	0.20 ± 0.03	480 ± 40
23	02/06–09/06	5,800 ± 400	0.25 ± 0.03	430 ± 30
24	09/06–16/06	5,100 ± 400	0.12 ± 0.03	250 ± 18
25	16/06–23/06	2,230 ± 160	< 0.10	141 ± 10

Continued on next page

Table A3 Continued

Week number	Period	^7Be $\mu\text{Bq}\cdot\text{m}^{-3}$	^{137}Cs $\mu\text{Bq}\cdot\text{m}^{-3}$	^{210}Pb $\mu\text{Bq}\cdot\text{m}^{-3}$
26	23/06–30/06	2,900 ± 200	0.07 ± 0.02	209 ± 15
27	30/06–07/07	1,680 ± 120	< 0.11	112 ± 8
28	07/07–14/07	2,700 ± 200	0.06 ± 0.02	191 ± 14
29	14/07–21/07	3,900 ± 300	0.12 ± 0.02	320 ± 20
30	21/07–28/07	3,200 ± 200	0.09 ± 0.02	450 ± 30
31	28/07–04/08	1,940 ± 140	< 0.09	130 ± 10
32	04/08–11/08	2,900 ± 200	< 0.09	147 ± 11
33	11/08–18/08	2,900 ± 200	0.10 ± 0.02	168 ± 12
34	18/08–25/08	4,800 ± 400	0.17 ± 0.03	350 ± 30
35	25/08–01/09	4,800 ± 400	0.24 ± 0.03	440 ± 30
36	01/09–08/09	2,900 ± 200	0.08 ± 0.02	224 ± 17
37	08/09–15/09	4,500 ± 300	0.23 ± 0.03	920 ± 70
38	15/09–22/09	5,100 ± 400	0.32 ± 0.03	620 ± 50
39	22/09–29/09	3,900 ± 300	0.21 ± 0.03	520 ± 40
40	29/09–06/10	3,000 ± 200	0.11 ± 0.03	194 ± 14
41	06/10–13/10	1,650 ± 120	0.23 ± 0.03	310 ± 20
42	13/10–20/10	2,610 ± 190	0.22 ± 0.03	680 ± 50
43	20/10–27/10	1,350 ± 100	0.26 ± 0.03	610 ± 40
44	27/10–03/11	3,300 ± 200	0.27 ± 0.03	280 ± 20
45	03/11–10/11	1,470 ± 110	0.18 ± 0.03	270 ± 20
46	10/11–17/11	1,720 ± 130	0.19 ± 0.03	430 ± 30
47	17/11–24/11	3,300 ± 200	0.12 ± 0.02	186 ± 14
48 ⁽¹⁾	24/11–30/11	2,250 ± 170	0.31 ± 0.04	220 ± 16
49 ⁽¹⁾	30/11–08/12	3,010 ± 170	0.54 ± 0.04	590 ± 30
50	08/12–15/12	2,310 ± 170	0.17 ± 0.03	830 ± 60
51	15/12–22/12	2,290 ± 170	0.36 ± 0.04	690 ± 50
52	22/12–29/12	1,290 ± 90	0.064 ± 0.019	97 ± 7
Average		2,910 ± 30 ⁽²⁾	0.208 ± 0.004 ^(2, 3)	312 ± 4 ⁽²⁾
SD ⁽⁴⁾		1,100	0.14	200

⁽¹⁾ The sampling period deviated from the regular 7-day sampling period.

⁽²⁾ The uncertainty in the yearly average is equal to the square root of the sum of the squared weekly uncertainties divided by the number of weeks. Uncertainties are given as 1σ .

⁽³⁾ The detection limits are omitted in the calculation of the averages.

⁽⁴⁾ SD is the standard deviation of the weekly results.

Table A4 Precipitation per month and monthly deposited ^3H , long-lived gross α and gross β activity sampled at RIVM in 2016

Month	Precipitation mm	^3H ⁽¹⁾ Bq·m ⁻²	Gross α Bq·m ⁻²	Gross β Bq·m ⁻²
January	129.5	< 200	3.6 ± 0.4	7.6 ± 0.6
February	91.0	< 160	2.3 ± 0.3	2.8 ± 0.2
March	44.8	< 80	3.9 ± 0.4	6.7 ± 0.5
April	71.5	< 110	7.6 ± 0.7	9.7 ± 0.7
May	59.5	< 90	7.0 ± 0.7	8.8 ± 0.7
June	141.7	< 200	7.8 ± 0.7	11.8 ± 0.9
July	93.0	< 170	7.3 ± 0.7	8.1 ± 0.6
August	56.1	< 100	6.4 ± 0.6	7.9 ± 0.6
September	16.2	32 ± 9	8.4 ± 0.8	9.6 ± 0.7
October	50.1	< 100	4.7 ± 0.5	5.1 ± 0.4
November	81.3	< 170	3.2 ± 0.4	5.2 ± 0.4
December	18.4	< 40	3.4 ± 0.4	4.8 ± 0.4
Total	852.8	-	66 ± 2 ⁽²⁾	88 ± 2 ⁽²⁾
Lower limit ⁽³⁾	-	22		
Upper limit ⁽³⁾	-	1,510		

⁽¹⁾ The detection limit in Bq·m⁻² is mainly dependent on the amount of precipitation, as the detection limit of the counting sample itself is more or less constant (1.6–2.0 Bq·L⁻¹).

⁽²⁾ The uncertainty in the sum is equal to the square root of the sum of the squared monthly uncertainties. Uncertainties are given as 1 σ .

⁽³⁾ The lower and upper limits are defined in Appendix B.

Table A5 Monthly deposited ^{210}Po activity ⁽¹⁾ sampled at RIVM in 2016

Month	^{210}Po Bq·m ⁻²
January	3.8 ± 0.3
February	1.08 ± 0.13
March	2.81 ± 0.16
April	5.9 ± 0.4
May	2.1 ± 0.3
June	3.5 ± 0.3
July	5.6 ± 0.4
August	4.7 ± 0.3
September	6.0 ± 0.3
October	2.8 ± 0.2
November ⁽³⁾	-
December	2.36 ± 0.13
Total	40.6 ± 1.0 ⁽²⁾
Lower limit ⁽⁴⁾	-
Upper limit ⁽⁴⁾	-

⁽¹⁾ Measurements were carried out using α -spectroscopy. Uncertainties are given as 1 σ .

⁽²⁾ The uncertainty in the sum is equal to the square root of the sum of the squared monthly uncertainties. Uncertainties are given as 1 σ .

⁽³⁾ The result for November has been rejected.

⁽⁴⁾ The lower and upper limits are defined in Appendix B.

Table A6 Yearly totals for long-lived gross α , gross β and ^3H activity in deposition since 1993

Year	Precipitation mm	^3H Bq·m ⁻²	Gross α Bq·m ⁻²	Gross β Bq·m ⁻²
1993	886	1,310 ± 30	54.3 ± 0.7	87.8 ± 0.8
1994	1,039	1,210 ± 30	52 ± 2	91 ± 3
1995	724	970 ± 40	33.6-44.6	95 ± 8
1996	626	970 ± 50	16.4 ± 1.5	67 ± 5
1997	760	1,160 ± 60	22.0-25.0	87 ± 3
1998	1,238	1,090-2,190	31.1 ± 1.3	106 ± 3
1999	916	1,420-1,900	25.5 ± 1.1	84 ± 2
2000	935	260-1,440	35.2 ± 1.3	104 ± 3
2001	1,053	0-2,420	23.9 ± 1	97 ± 3
2002	965	300-1,710	20.6 ± 0.9	97 ± 2
2003	605	260-1,080	13.6-16.7	70.0 ± 1.8
2004	875	0-1,600	14.3-17.1	73.5 ± 1.8
2005	856	0-1,530	17.6 ± 1.0	88 ± 2
2006	854	280-1,820	25.7 ± 1.5	98 ± 3
2007	984	335-1,600	24.4 ± 1.2	85 ± 2
2008	908	102-1,550	39.4 ± 1.5	106 ± 3
2009	794	0-1,330	36.9 ± 1.3	95 ± 2
2010	868	180-1,400	36.7 ± 1.3	90 ± 2
2011	895	332-1,540	45.0 ± 1.5	123 ± 3
2012	922	316-1,650	32.7 ± 1.1	88 ± 2
2013	851	0-1,580	48.1 ± 1.6	95 ± 2
2014	878	117-1,630	62 ± 2	106 ± 3
2015	878	210-1,650	53.9 ± 1.8	90 ± 2
2016	853	22-1,510	66 ± 2	88 ± 2

Either the yearly total with uncertainty ⁽¹⁾ or the lower and upper limits ⁽²⁾ of the 68% confidence interval are given.

⁽¹⁾ Uncertainties are given as 1σ .

⁽²⁾ Lower and upper limits are given as defined in Appendix B.

Table A7 Yearly totals for ^7Be , ^{137}Cs , ^{210}Pb and ^{210}Po activity in deposition since 1993

Year	^7Be ⁽³⁾ Bq·m ⁻²	^{137}Cs ⁽³⁾ Bq·m ⁻²	^{210}Pb ⁽³⁾ Bq·m ⁻²	^{210}Pb ⁽⁴⁾ Bq·m ⁻²	^{210}Po ⁽⁴⁾ Bq·m ⁻²
1993	1,090 ± 20	0.50–0.76	105 ± 2	78 ± 3	7.2 ± 0.5
1994	1,320 ± 30	0.36–0.71	118 ± 3	82 ± 3	12.0–14.2
1995	990 ± 20	0.37–0.63	96 ± 2	n/a ⁽⁵⁾	n/a ⁽⁵⁾
1996	920 ± 20	0.52–0.83	63–67	57 ± 3	9 ± 2
1997	1,090 ± 30	0.11–0.69	65–69	80 ± 4	0–10.2
1998	1,840 ± 50	0.56–0.85	162 ± 4	91 ± 4	3.0–15.1
1999	1,580 ± 30	1.16–1.99	158 ± 4	- ⁽⁶⁾	0.7–5.3
2000	1,490 ± 30	0–4.82	177 ± 6	-	0.6–8.0
2001	1,480 ± 30	0–4.50	83–104	-	6.5–9.4
2002	1,510 ± 30	0–5.22	119–142	-	6.1–8.5
2003	1,000–1,050	0–4.69	88–113	-	4.3–5.6
2004	1,330 ± 30	0.22–5.53	64–102	-	5.4–7.7
2005	1,320 ± 30	0–6.09	87–117	-	8.9–10.2
2006	1,400 ± 30	0.06–7.47	66–103	-	14.8–16.4 ⁽⁷⁾
2007	1,760 ± 40	0.11–7.37	72–132	-	13.4 ± 0.4 ⁽⁷⁾
2008	1,990 ± 40	0–7.63	63–143	-	29.4 ± 0.7
2009	1,410 ± 30	0–4.3	82–125	-	32.5 ± 0.7
2010	1,240 ± 30	0–1.2	93 ± 2	-	33.2 ± 0.8
2011	1,320 ± 30	0.5–1.5	104 ± 2	-	61.4 ± 1.0
2012	1,330 ± 30	0–1.2	98 ± 2	-	33.8 ± 0.6
2013	1,030 ± 30	0–1.1	82.9 ± 1.8	-	21.2 ± 0.6 ⁽⁸⁾
2014	1,341 ± 17	0–1.1	107 ± 2	-	41.4 ± 1.0
2015	1,219 ± 16	0–1.1	102.8 ± 1.5	-	40.7 ± 1.0
2016	1,375 ± 18	0.04–1.15	95 – 98	-	40.6 ± 1.0 ⁽⁹⁾

Either the yearly total with uncertainty ⁽¹⁾ or the lower and upper limits ⁽²⁾ of the 68% confidence range are given.

⁽¹⁾ Uncertainties are given as 1σ .

⁽²⁾ Lower and upper limits are given as defined in Appendix B.

⁽³⁾ Data from γ -spectroscopy.

⁽⁴⁾ Data from α -spectroscopy.

⁽⁵⁾ Not available. Result rejected [75].

⁽⁶⁾ α -spectroscopy analysis of ^{210}Pb stopped in 1999.

⁽⁷⁾ Results revised in RIVM Report 610791003.

⁽⁸⁾ The yearly total deposition is based on 10 monthly results.

⁽⁹⁾ The yearly total deposition is based on 11 monthly results.

Table A8 Weekly deposited ^7Be , ^{137}Cs and ^{210}Pb activity ⁽¹⁾ sampled at RIVM in 2016

Week no.	Period	Precipitation mm	^7Be Bq·m ⁻²	^{137}Cs Bq·m ⁻²	^{210}Pb Bq·m ⁻²
1	31/12-07/01	27.5	8.7 ± 0.6	< 0.01	0.61 ± 0.06
2	07/01-14/01	34.8	33 ± 2	< 0.02	0.83 ± 0.10
3	14/01-21/01	14.3	20.8 ± 1.5	< 0.02	0.56 ± 0.08
4	21/01-28/01	23.5	20.4 ± 1.5	< 0.02	0.91 ± 0.10
5	28/01-04/02	29.5	50 ± 4	< 0.02	1.73 ± 0.16
6	04/02-11/02	32.5	38 ± 3	< 0.02	1.41 ± 0.13
7	11/02-18/02	18.0	12.1 ± 0.9	< 0.02	2.03 ± 0.17
8	18/02-25/02	25.0	24.7 ± 1.8	< 0.02	0.99 ± 0.11
9	25/02-03/03	15.5	19.4 ± 1.4	< 0.02	0.71 ± 0.10
10	03/03-10/03	18.5	18.2 ± 1.3	< 0.02	1.21 ± 0.13
11	10/03-17/03	0.0	1.52 ± 0.14	< 0.02	0.24 ± 0.07
12	17/03-24/03	2.3	6.2 ± 0.5	< 0.03	0.63 ± 0.10
13	24/03-31/03	24.0	38 ± 3	< 0.02	1.41 ± 0.13
14	31/03-07/04	26.0	60 ± 4	< 0.02	5.4 ± 0.4
15	07/04-14/04	9.5	22.2 ± 1.6	< 0.02	0.67 ± 0.09
16	14/04-21/04	13.0	24.7 ± 1.8	< 0.02	1.05 ± 0.11
17	21/04-28/04	23.0	49 ± 4	< 0.02	2.6 ± 0.2
18	28/04-04/05	14.5	27 ± 2	< 0.02	1.38 ± 0.11
19	04/05-12/05	0.0	6.8 ± 0.5	0.015 ± 0.005	3.1 ± 0.2
20	12/05-19/05	1.5	23.8 ± 1.7	0.031 ± 0.005	6.8 ± 0.5
21	19/05-26/05	35.5	46 ± 3	< 0.02	3.3 ± 0.3
22	26/05-02/06	8.0	30 ± 2	< 0.02	3.5 ± 0.3
23	02/06-09/06	1.7	20.9 ± 1.5	< 0.02	2.04 ± 0.17
24	09/06-16/06	28.5	46 ± 3	< 0.02	3.0 ± 0.2
25	16/06-23/06	72.5	93 ± 7	< 0.03	4.7 ± 0.4
26	23/06-30/06	39.0	85 ± 6	< 0.02	3.9 ± 0.3

Continued on next page

Table A8 Continued

Week no.	Period	Precipitation mm	⁷ Be Bq·m ⁻²	¹³⁷ Cs Bq·m ⁻²	²¹⁰ Pb Bq·m ⁻²
27	30/06-07/07	27.5	42 ± 3	< 0.02	1.73 ± 0.16
28	07/07-14/07	15.0	34 ± 2	< 0.03	2.5 ± 0.2
29	14/07-21/07	0.0	10.9 ± 0.8	< 0.02	3.2 ± 0.2
30	21/07-28/07	22.5	58 ± 4	< 0.02	5.5 ± 0.4
31	28/07-04/08	28.0	37 ± 3	< 0.02	1.32 ± 0.13
32	04/08-11/08	24.0	48 ± 4	< 0.02	2.4 ± 0.2
33	11/08-18/08	10.0	11.7 ± 0.9	< 0.02	0.43 ± 0.08
34	18/08-25/08	21.0	37 ± 3	< 0.02	1.85 ± 0.16
35	25/08-01/09	1.1	28 ± 2	< 0.02	4.2 ± 0.3
36	01/09-08/09	3.9	17.9 ± 1.4	< 0.02	1.41 ± 0.14
37	08/09-15/09	1.5	13.5 ± 1.0	< 0.02	1.60 ± 0.15
38	15/09-22/09	6.3	18.2 ± 1.4	< 0.02	1.52 ± 0.14
39	22/09-29/09	4.5	16.9 ± 1.3	< 0.02	2.23 ± 0.19
40	29/09-06/10	12.0	14.4 ± 1.1	< 0.02	1.08 ± 0.11
41	06/10-13/10	0.0	3.7 ± 0.3	< 0.02	0.60 ± 0.09
42	13/10-20/10	31.0	30 ± 2	< 0.02	2.8 ± 0.2
43	20/10-27/10	1.7	4.4 ± 0.3	< 0.02	0.18 ± 0.07
44	27/10-03/11	5.4	7.9 ± 0.6	< 0.02	0.91 ± 0.11
45	03/11-10/11	24.5	23.6 ± 1.7	< 0.02	0.77 ± 0.10
46	10/11-17/11	32.5	38 ± 3	< 0.02	0.82 ± 0.11
47	17/11-24/11	23.8	23.4 ± 1.8	< 0.018	1.25 ± 0.12
48	24/11-30/11	0.5	4.6 ± 0.4	< 0.02	0.36 ± 0.07
49	30/11-08/12	1.9	2.7 ± 0.2	< 0.02	< 0.2
50	08/12-15/12	8.0	9.3 ± 0.7	< 0.02	1.30 ± 0.13
51	15/12-22/12	3.9	7.1 ± 0.5	< 0.02	0.79 ± 0.10
52	22/12-29/12	4.6	6.5 ± 0.5	< 0.02	0.39 ± 0.09
Total ⁽²⁾		852.8	1,375 ± 18	-	-
Lower limit ⁽³⁾		-	-	0.04	95
Upper limit ⁽³⁾		-	-	1.15	98

⁽¹⁾ Measurements were carried out using γ -spectroscopy.

⁽²⁾ The uncertainty in the sum is equal to the square root of the sum of the squared weekly uncertainties. Uncertainties are given as 1σ .

⁽³⁾ The lower and upper limits are defined in Appendix B.

Table A9 Yearly average α activity concentration in air and ambient dose equivalent rate in 2016, as measured by the NMR stations equipped with aerosol monitors

Station	No.	α activity concentration Bq.m⁻³	Ambient dose equivalent rate ⁽¹⁾ nSv.h⁻¹
Arnhem	970	4.1	77
Kollumerwaard	972	3.4	87
Valthermond	974	3.0	68
Vlaardingen	976	3.2	79
Braakman	978	3.7	75
Huijbergen	980	3.5	64
Houtakker	982	3.7	71
Wijnandsrade	984	7.1	86
Eibergen	986	4.2	68
De Zilk	988	3.3	72
Wieringerwerf	990	2.5	81
Vredepeel	992	4.0	63
Biddinghuizen	994	3.5	87
Bilthoven	998	3.2	68

⁽¹⁾ These dose equivalent rate monitors are placed differently from the dose equivalent rate monitors mentioned in Table A10 with regard to height and surface covering.

Table A10 Yearly average ambient dose equivalent rate for the NMR stations in 2017

Station	No.	Ambient dose equivalent rate nSv.h ⁻¹	Station	No.	Ambient dose equivalent rate nSv.h ⁻¹
Den Burg	1001	77	Lelystad	1103	82
Den Helder	1002	76	Urk	1105	82
Den Oever	1003	78	Eemshaven	1106	82
Petten	1006	70	Uithuizen	1107	87
Kolhorn	1007	92	Wagenborgen	1109	78
Egmond aan Zee	1009	70	Winschoten	1110	81
Heerhugowaard	1011	84	Ter Apel	1111	74
Nederhorst Den Berg	1015	82	Stadskanaal	1112	72
Velsen	1016	75	Nieuweschans	1113	84
Enkhuizen	1018	81	Bellingwolde	1114	66
Oosthuizen	1019	80	Groningen	1116	85
Zaandam ⁽¹⁾	1021	111	Leens	1117	87
Gouda	1024	81	Grijpskerk	1118	79
Dordrecht	1027	68	Meppel	1125	76
Zuid Beijerland	1028	87	Hoogeveen	1126	70
Rotterdam-Schiebroek	1031	73	Steenwijksmoer	1129	73
Pijnacker	1032	83	Nw. Amsterdam	1130	81
Maasvlakte	1035	83	Nw. Schoonebeek/	1131	70
Rotterdam-Waalhaven	1036	65	Weiteveen		
Maassluis	1037	98	Emmen	1132	80
Hellevoetsluis	1038	112	Laren (Gld) ⁽²⁾	1134	73
Ouddorp	1039	70	Hengelo (Ov) ⁽²⁾	1135	79
Otterlo ⁽²⁾	1042	87	Vroomshoop ⁽²⁾	1138	76
Wageningen	1043	74	Enschede	1139	74
Amersfoort	1046	76	Losser	1140	69
Harderwijk	1050	71	Oldenzaal	1141	78
Wijk bij Duurstede	1056	90	Rijssen	1143	83
Nieuwegein	1062	90	's Heerenberg	1144	81
Zegveld	1063	74	Dinxperlo	1145	90
Lopik (Cabauw)	1064	86	Varsseveld	1146	81
Apeldoorn	1066	77	Groenlo	1147	93
Heerenveen	1071	69	Deventer	1148	87
Oosterwolde	1072	87	Etten-Leur	1154	75
Bergum	1074	75	Den Bosch	1157	77
Witmarsum ⁽²⁾	1076	99	Raamsdonkveer	1159	94
Sneek	1077	78	Ulvenhout	1160	73
St Jacobiparochie	1081	86	Baarle Nassau	1161	80
Holwerd	1082	92	Mill	1163	73
Leeuwarden	1085	79	Volkel	1164	67
Zwolle	1088	86	Oss	1167	78
Ommen	1093	71	Nuenen	1172	77
Hardenberg	1095	74	Bergeijk	1174	100
Assen	1097	74	Waalre	1175	76
Rutten	1099	85	Someren (Dorp)	1176	75

Continued on next page

Table A10 Continued

Station	No.	Ambient dose equivalent rate nSv.h ⁻¹	Station	No.	Ambient dose equivalent rate nSv.h ⁻¹
Oisterwijk	1178	79	Hoensbroek	1225	96
Riel	1179	80	Gennep	1228	75
Oostelbeers	1180	110	Elst (Gld)	1229	84
Hilvarenbeek	1181	74	Zevenaar	1230	82
Venray	1183	66	Nijmegen	1231	75
Nieuw-Bergen	1184	66	Amstelveen	1233	88
Sevenum	1185	80	Amsterdam	1234	75
Reuver	1188	76	Aalsmeer	1236	82
Nederweert	1189	79	Nispen	1237	71
Heythuysen	1190	85	Groesbeek	1240	80
Mariahoop	1191	80	Tubbergen	1243	78
Stramproy	1192	71	Haaksbergen	1244	72
Eerbeek	1193	79	Scheveningen	1247	84
Leiden	1196	82	Zaltbommel	1251	82
Hulst	1197	85	IJzendijke	1252	84
Terneuzen	1199	81	Ritthem	1253	103
Sluis	1201	82	Vlissingen Haven	1254	80
Vlissingen	1202	87	Nieuwdorp	1255	82
Halsteren	1204	74	's-Heerenhoek ⁽³⁾	1256	128
Oud Gastel	1206	76	Driewegen	1257	95
Goes	1207	80	Arnhem	1258	85
Bruinisse	1209	82	Heinkenszand	1259	98
Burgh- Haamstede	1211	67	Baarland	1260	93
Vrouwenpolder	1212	68	Biervliet	1261	77
Yerseke	1213	85	Nummer Een	1262	89
Middelburg	1215	88	Rilland	1263	84
Westkapelle	1216	77	Putte	1264	63
Maasband	1218	81	Nieuw Namen	1265	93
Maastricht	1220	104	Beneden Leeuwen	1272	87
Ravensbos (Arensgehout)	1221	95	Denekamp	1278	73
Vaals	1222	95	Winterswijk (Kotten)	1279	75
Gulpen	1223	84	Bilthoven	1280	62
Kerkrade	1224	94	Gastel (Maarheze)	1281	80

⁽¹⁾ The Zaandam station showed a significantly higher value than most other stations. This is due to a higher background level of the surrounding surface at the site since the end of 2014.

⁽²⁾ A few stations have been moved during 2018: Wekerom station was moved to Otterlo, Harlingen station to Witmarsum, Hengelo (Ov) station to Laren (Gld) and Den Ham station to Vroomshoop.

⁽³⁾ As in previous years, the 's-Heerenhoek station showed a significantly higher value than all other stations. This is due to a higher background level of the ground surface at the site. Since September 2009, this background level has been reduced by covering the surrounding ground surface with a layer of shells.

Table A11 Gross α , residual β , ^3H , ^{90}Sr and ^{226}Ra activity concentrations ($\text{mBq}\cdot\text{L}^{-1}$) in surface water in 2016 as measured by RWS

Date	Gross α $\text{mBq}\cdot\text{L}^{-1}$	Residual β $\text{mBq}\cdot\text{L}^{-1}$	^3H $\text{mBq}\cdot\text{L}^{-1}$	^{90}Sr $\text{mBq}\cdot\text{L}^{-1}$	^{226}Ra $\text{mBq}\cdot\text{L}^{-1}$
Location IJsselmeer					
26/01/16	59	31	2,210		
24/02/16	53	21			
23/03/16	44	23	2,280		
17/05/16	34	3	2,180		
15/06/16	27	10			
13/07/16	40	< 1	2,700		
10/08/16	29	18			
07/09/16	2	4	2,640		
05/10/16	34	15			
02/11/16	23	5	2,550		
01/12/16	17	< 1			
27/12/16	33	41	2,310		
Average	33	14	2,410		
Location Noordzeekanaal					
11/01/16	182	5	3,030		
08/02/16	24	14	2,980		
07/03/16	61	15	2,450		
04/04/16	121	23	2,760		
02/05/16	40	10	2,580		
30/05/16	78	55	2,390		
27/06/16	72	13	2,490		
25/07/16	53	7	2,820		
22/08/16	198	18	2,040		
19/09/16	91	23	2,400		
17/10/16	176	51	2,330		
14/11/16	162	36	2,310		
12/12/16	156	5	2,740		
Average	109	21	2,560		
Location Nieuwe Waterweg					
28/01/16	67	87			
25/02/16	101	107	2,860	< 1	6.4
22/03/16	83	48			
21/04/16	58	53	3,430	< 1	2.6
19/05/16	43	43			
13/06/16	14	45	1,940	2.0	3.9
14/07/16	46	13			
11/08/16	78	31	3,080	< 1	3.3
08/09/16	150	33			
06/10/16	100	19	3,380	< 1	3.1
03/11/16	245	18			
01/12/16	229	86	4,390	< 1	4.7
29/12/16	36	17			
Average	96	46	3180	< 1	4.0

Continued on next page

Table A11 Continued

Date	Gross α mBq·L ⁻¹	Residual β mBq·L ⁻¹	³ H mBq·L ⁻¹	⁹⁰ Sr mBq·L ⁻¹	²²⁶ Ra mBq·L ⁻¹
Location	Rhine				
06/01/16	45	33	3,420	< 1	27.6
03/02/16	70	41	1,550		
02/03/16	41	33	4,570	< 1	3.2
30/03/16	34	11	4,070		
28/04/16	58	32	3,160	2.5	7.4
25/05/16	46	23	3,430		
22/06/16	53	47	2,870	< 1	2.9
20/07/16	58	37	2,630		
17/08/16	38	24	3,250	4.1	4.8
14/09/16	60	26	2,920		
12/10/16	49	31	4,480	4.1	10.4
09/11/16	33	26	3,920		
07/12/16	54	16	5,180	9.3	3.5
Average	49	29	3,500	3.1	8.6
Location	Scheldt				
27/01/16	170	212	6,660		16.7
24/02/16	105	91			
21/03/16	86	71	5,620		3.9
17/05/16	180	110	11,300		61.8
14/06/16	97	78			
14/07/16	150	33	4,210		3.2
08/08/16	138	63			
06/09/16	369	108	16,400		14.9
04/10/16	392	165			
02/11/16	290	171	16,400		8.4
01/12/16	648	139			
29/12/16	364	103	14,900		123.0
Average	249	112	10,800		33.1
Location	Meuse				
05/01/16	49	35	1,160	< 1	9.0
02/02/16	101	118	868		
01/03/16	18	25	11,500	< 1	1.8
29/03/16	33	18	1,050		
26/04/16	14	10	788	< 1	5.1
24/05/16	37	14	1,980		
21/06/16	43	40	8,050	5.3	1.3
19/07/16	47	4	4,620		
16/08/16	25	12	33,800	< 1	5.6
13/09/16	19	19	23,400		
11/10/16	12	5	16,500	1.4	3.6
08/11/16	28	< 1	55,900		
06/12/16	19	11	29,600	< 1	12.0
Average	34	24	14,600	< 1.3	5.5

Continued on next page

Table A11 Continued

Date	Gross α mBq·L⁻¹	Residual β mBq·L⁻¹	³H mBq·L⁻¹	⁹⁰Sr mBq·L⁻¹	²²⁶Ra mBq·L⁻¹
Location	Kanaal Gent Terneuzen				
20/01/16	60	28			
16/02/16	82	62	1,500		
14/03/16	58	14			
13/04/16	74	22	920		
12/05/16	45	41			
07/06/16	60	23	1,160		
05/07/16	43	17			
01/08/16	90	27	1,340		
29/08/16	76	18			
28/09/16	127	37	2,620		
25/10/16	115	19			
23/11/16	18	35	1,760		
20/12/16	132	20			
Average	75	28	1,550		
Location	Haringvliet				
21/01/16	30	23	7,920		
18/02/16	38	31			
17/03/16	13	7	6,420		
14/04/16	15	19			
12/05/16	26	3	4,710		
09/06/16	36	19			
07/07/16	44	11	2,740		
04/08/16	44	8			
01/09/16	28	16	4,580		
29/09/16	26	2			
27/10/16	20	< 1	4,030		
24/11/16	31	26			
22/12/16	29	1	4,700		
Average	29	13	5,010		

Table A12 ^{60}Co , ^{131}I , ^{137}Cs and ^{210}Pb activity concentrations in suspended solids ($\text{Bq}\cdot\text{kg}^{-1}$) in surface water in 2015 as measured by RWS

Date	^{60}Co $\text{Bq}\cdot\text{kg}^{-1}$	^{131}I $\text{Bq}\cdot\text{kg}^{-1}$	^{137}Cs $\text{Bq}\cdot\text{kg}^{-1}$	^{210}Pb $\text{Bq}\cdot\text{kg}^{-1}$
Location	IJsselmeer			
26/01/16	< 1	< 1	4.6	
24/02/16	< 1	< 1	5.9	
23/03/16	< 1	< 1	5.2	
20/04/16	< 1	< 1	4.8	
17/05/16	< 1	< 1	4.4	
15/06/16	< 1	< 1	4.9	
13/07/16	< 1	< 1	< 1	
10/08/16	< 1	< 1	1.3	
07/09/16	< 1	< 1	< 1	
05/10/16	< 1	< 1	2.1	
02/11/16	< 1	< 1	2.3	
01/12/16	< 1	< 1	1.2	
27/12/16	< 1	< 1	2.0	
Average	< 1	< 1	3.0	
Location	Noordzeekanaal			
11/01/16	< 1	8.7	7.8	
07/03/16	< 1	8.0	9.3	
02/05/16	< 1	25	6.2	
27/06/16	< 1	< 1	6.7	
22/08/16	< 1	104	2.3	
18/10/16	< 1	63	4.9	
12/12/16	< 1	43	6.5	
Average	< 1	36	6.2	
Location	Nieuwe Waterweg			
14/01/16	< 1	< 1	6.1	
11/02/16	< 1	2.2	6.1	87
10/03/16	< 1	< 1	6.7	
07/04/16	< 1	< 1	6.5	102
03/05/16	< 1	< 1	9.1	
02/06/16	< 1	4.0	8.8	125
30/06/16	< 1	7.9	9.2	
28/07/16	< 1	< 1	8.5	101
25/08/16	1.8	< 1	8.3	
22/09/16	< 1	< 1	6.5	81
20/10/16	< 1	< 1	5.0	
17/11/16	< 1	< 1	6.3	85
15/12/16	< 1	6.1	8.4	
Average	< 1	< 2	7.3	97
Location	Rhine			
06/01/16	< 1	< 1	12.4	131
03/02/16	< 1	< 1	11.6	
17/02/16	< 1	6.2	13.4	
02/03/16	< 1	8.1	11.9	119

Continued on next page

Table A12 Continued

Date	⁶⁰ Co Bq·kg ⁻¹	¹³¹ I Bq·kg ⁻¹	¹³⁷ Cs Bq·kg ⁻¹	²¹⁰ Pb Bq·kg ⁻¹
Location	Rhine			
16/03/16	< 1	< 1	13.3	
30/03/16	< 1	< 1	9.0	
13/04/16	< 1	< 1	9.5	
28/04/16	< 1	5.0	12.2	135
11/05/16	< 1	6.7	10.2	
25/05/16	< 1	< 1	9.1	
08/06/16	< 1	< 1	11.7	
22/06/16	< 1	4.0	11.8	109
06/07/16	< 1	< 1	10.3	
20/07/16	< 1	< 1	10.6	
03/08/16	< 1	< 1	9.1	
17/08/16	< 1	< 1	8.4	116
31/08/16	< 1	< 1	8.7	
14/09/16	< 1	< 1	9.4	
28/09/16	< 1	< 1	7.8	
12/10/16	< 1	< 1	8.7	118
26/10/16	< 1	< 1	10.1	
09/11/16	< 1	< 1	9.3	
23/11/16	< 1	11.0	10.0	
07/12/16	< 1	< 1	10.7	121
21/12/16	< 1	< 1	9.0	
Average	< 1	< 2	10.3	121
Location	Scheldt			
27/01/16	< 1	3.6	5.6	78
24/02/16	< 1	< 1	5.6	
21/03/16	< 1	< 1	6.5	86
22/04/16	< 1	< 1	6.5	
17/05/16	< 1	< 1	5.4	79
14/06/16	< 1	1.8	6.1	
14/07/16	< 1	< 1	6.4	86
08/08/16	< 1	< 1	5.7	
06/09/16	< 1	< 1	5.8	92
04/10/16	< 1	< 1	4.4	
02/11/16	< 1	< 1	5.8	73
01/12/16	< 1	< 1	5.1	
29/12/16	< 1	< 1	4.9	68
Average	< 1	< 1	5.7	80
Location	Meuse			
04/01/16	9.9	11.3	11.3	155
11/01/16	1.6	3.5	12.2	
18/01/16	14.4	7.1	16.3	
25/01/16	< 1	12.2	11.5	
01/02/16	< 1	5.0	9.3	

Continued on next page

Table A12 Continued

Date	⁶⁰ Co Bq·kg ⁻¹	¹³¹ I Bq·kg ⁻¹	¹³⁷ Cs Bq·kg ⁻¹	²¹⁰ Pb Bq·kg ⁻¹
Location	Meuse			
08/02/16	< 1	6.0	9.3	
15/02/16	5.7	< 1	10.1	
22/02/16	< 1	4.8	10.0	
29/02/16	< 1	7.1	11.1	112
07/03/16	< 1	5.7	9.8	
14/03/16	14.4	4.5	10.7	
21/03/16	95	12.4	9.3	
29/03/16	17.3	13.8	8.7	
04/04/16	6.5	7.5	10.6	
11/04/16	37.4	10.4	9.9	
18/04/16	15.2	< 1	8.3	
25/04/16	10.6	< 1	8.3	129
02/05/16	30.4	18.1	9.0	
09/05/16	12.2	4.3	7.5	
17/05/16	41	22.6	8.3	
23/05/16	24.4	15.5	9.0	
30/05/16	8.6	< 1	7.8	
06/06/16	< 1	6.5	11.9	
13/06/16	3.4	< 1	9.9	
20/06/16	2.3	7.2	10.7	123
27/06/16	7.3	12.5	10.8	
04/07/16	9.0	< 1	11.8	
11/07/16	5.5	11.3	10.5	
18/07/16	7.0	11.1	10.3	
25/07/16	5.5	3.2	10.3	
02/08/16	8.9	< 1	10.7	
08/08/16	9.3	< 1	10.6	
16/08/16	10.3	< 1	12.0	141
22/08/16	7.5	< 1	10.6	
29/08/16	5.8	< 1	8.4	
05/09/16	4.3	< 1	5.3	
12/09/16	3.8	< 1	4.7	
20/09/16	3.3	25	6.9	
26/09/16	4.7	12.8	6.9	
10/10/16	3.8	< 1	7.9	138
17/10/16	5.6	< 1	12.5	
25/10/16	3.8	< 1	12.6	
31/10/16	5.7	< 1	12.7	
07/11/16	4.2	< 1	12.1	
14/11/16	6.3	< 1	9.7	
21/11/16	6.5	11.2	10.7	
28/11/16	9.6	18.8	12.7	
05/12/16	12.7	< 1	11.5	185
12/12/16	6.1	< 1	12	
19/12/16	7.7	16.0	10.8	
27/12/16	13.2	< 1	12	
Average	10.4	6.2	10.1	140

Continued on next page

Table A12 Continued

Date	⁶⁰Co Bq·kg⁻¹	¹³¹I Bq·kg⁻¹	¹³⁷Cs Bq·kg⁻¹	²¹⁰Pb Bq·kg⁻¹
Location	Kanaal Gent Terneuzen			
25/02/16	< 1	9.2	5.8	
25/05/16	< 1	< 1	3.1	
24/08/16	< 1	< 1	3.8	
23/11/16	< 1	15.8	8.3	
Average	< 1	6.5	5.3	
Location	Haringvliet			
21/01/16	< 1	6.7	12.8	
18/02/16	< 1	< 1	14.6	
17/03/16	< 1	< 1	13.7	
14/04/16	< 1	< 1	11.1	
12/05/16	< 1	< 1	7.3	
09/06/16	< 1	< 1	12.6	
07/07/16	< 1	< 1	13.5	
04/08/16	< 1	< 1	13.4	
24/11/16	< 1	< 1	12.5	
Average	< 1	< 1.2	12.4	

Table A13 Gross α , residual β , ^3H and ^{90}Sr activity concentrations ($\text{mBq}\cdot\text{L}^{-1}$) in seawater in 2016 as measured by RWS

Date	Gross α $\text{mBq}\cdot\text{L}^{-1}$	Residual β $\text{mBq}\cdot\text{L}^{-1}$	^3H $\text{mBq}\cdot\text{L}^{-1}$	^{90}Sr $\text{mBq}\cdot\text{L}^{-1}$
Location Coastal Area				
15/02/16	424	110	5,770	
17/05/16	183	67	4,220	
15/08/16	451	16	3,410	
14/11/16	563	43	3,730	
Average	410	59	4,300	
Location Southern North Sea				
15/02/16	195	43	4,720	< 1
17/05/16	104	62	824	< 1
15/08/16	543	31	1,850	2.3
14/11/16	548	37	3,090	4.4
Average	350	43	2,600	1.9
Location Central North Sea				
12/01/16	145	34	293	< 1
12/04/16	163	37	272	< 1
14/06/16	209	29	615	< 1
17/08/16	229	34	224	4.1
Average	190	34	350	< 1.4
Location Delta Coastal Waters				
13/01/16	40	76	6,320	
17/02/16	906	51	4,900	< 1
16/03/16	166	54	5,650	
14/04/16	169	31	5,330	
19/05/16	299	74	4,090	< 1
16/06/16	183	38	4,070	
14/07/16	216	27	4,170	
18/08/16	344	27	3,770	< 1
14/09/16	362	35	4,460	
12/10/16	292	15	4,330	
16/11/16	513	51	3,840	6.1
14/12/16	533	43	4,870	
Average	340	44	4,700	< 1.9
Location Western Scheldt				
11/01/16	222	283	6,350	2.2
08/02/16	306	285	7,000	2.9
07/03/16	801	74	6,270	< 1
06/04/16	289	124	5,150	< 1
03/05/16	324	72	5,740	< 1
30/05/16	282	111	4,780	< 1
29/06/16	228	40	4,470	2.4
25/07/16	380	24	4,340	4.1
22/08/16	358	34	3,830	< 1
20/09/16	372	53	4,270	< 1
18/10/16	298	106	4,670	1.5
17/11/16	261	102	4,940	1.7
12/12/16	418	59	5,900	5.3
Average	350	110	5,200	< 1.8

Continued on next page

Table A13 Continued

Location Eems-Dollard			
18/02/16	31	43	5,920
19/05/16	122	54	4,190
15/08/16	338	31	2,440
10/11/16	349	44	2,550
Average	210	43	3,800
Location Wadden Sea West			
10/02/16	210	90	6,070
19/05/16	494	61	3,680
16/08/16	287	63	3,260
16/11/16	563	58	3,410
Average	390	68	4,100
Location Wadden Sea East			
17/02/16	255	196	6,470
12/05/16	400	174	3,430
11/08/16	491	97	2,430
15/11/16	548	100	2,820
Average	420	140	3,800

Table A14 ^{137}Cs and ^{210}Pb activity concentrations in suspended solids ($\text{Bq}\cdot\text{kg}^{-1}$) in seawater in 2016 as measured by RWS

Date	^{137}Cs $\text{Bq}\cdot\text{kg}^{-1}$	^{210}Pb $\text{Bq}\cdot\text{kg}^{-1}$
Location Western Scheldt		
29/02/16	4.1	68
24/05/16	2.5	53
25/08/16	3.1	58
24/11/16	3.6	71
Average	3.3	62

Table A15 Monthly average gross α activity concentrations in air dust near the Borssele nuclear power plant in 2016

Date ⁽¹⁾	Gross α ⁽²⁾ $\text{mBq}\cdot\text{m}^{-3}$				
Location	21	22	23	27	29
03/02/16	0.021	0.011	0.046	0.03	0.002
02/03/16	0.022	0.014	0.005	0.008	0.013
06/04/16	0.008	0.026	0.025	0.01	0.006
03/05/16	0.018	0.017	0.051	0.026	0.003
07/06/16	0.095	0.04	0.053	0.015	0.003
06/07/16	0.021	0.028	0.005	0.005	0.005
03/08/16	0.031	0.026	0.015	0.032	0.006
07/09/16	0.021	0.055	0.012	0.035	0.066
06/10/16	0.059	0.069	0.041	0.054	0.031
03/11/16	0.104	0.029	0.02	0.051	-
05/12/16	0.064	0.043	0.013	0.028	0.01
05/01/17	0.06	0.065	0.153	0.05	0.051

⁽¹⁾ End date of monthly sampling period.

⁽²⁾ Gross α activity concentrations in air dust are given as indicative values.

Table A16 Monthly average gross β activity concentrations in air dust near the Borssele nuclear power plant in 2016

Date ⁽¹⁾	Gross β mBq·m ⁻³				
Location	21	22	23	27	29
03/02/16	0.03 ± 0.05	0.14 ± 0.03	0.27 ± 0.04	0.27 ± 0.05	0.03 ± 0.04
02/03/16	0.17 ± 0.05	0.22 ± 0.03	0.36 ± 0.04	0.06 ± 0.05	0.12 ± 0.04
06/04/16	0.31 ± 0.04	0.28 ± 0.02	0.39 ± 0.03	0.19 ± 0.04	0.08 ± 0.03
03/05/16	0.13 ± 0.05	0.19 ± 0.03	0.27 ± 0.04	0.14 ± 0.05	0.08 ± 0.05
07/06/16	0.65 ± 0.05	0.49 ± 0.02	0.34 ± 0.03	0.29 ± 0.04	0.10 ± 0.04
06/07/16	0.15 ± 0.05	0.33 ± 0.03	0.09 ± 0.04	0.11 ± 0.06	0.06 ± 0.02
03/08/16	0.33 ± 0.05	0.24 ± 0.03	0.06 ± 0.03	0.17 ± 0.04	0.21 ± 0.11
07/09/16	0.45 ± 0.05	0.62 ± 0.04	0.05 ± 0.03	0.33 ± 0.04	0.45 ± 0.08
06/10/16	1.01 ± 0.06	0.63 ± 0.03	0.07 ± 0.06	0.51 ± 0.05	0.45 ± 0.08
03/11/16	0.76 ± 0.07	0.41 ± 0.04	0.47 ± 0.07	0.53 ± 0.05	-
05/12/16	0.50 ± 0.05	0.43 ± 0.03	0.23 ± 0.06	0.10 ± 0.04	0.21 ± 0.03
05/01/17	1.12 ± 0.07	0.70 ± 0.04	1.27 ± 0.06	0.56 ± 0.05	0.73 ± 0.03

⁽¹⁾ End date of monthly sampling period.

Table A17 Monthly average activity concentrations of γ -emitters in air (dust) near the Borssele nuclear power plant in 2016 ⁽¹⁾

Date ⁽²⁾	⁶⁰ Co mBq·m ⁻³	¹³¹ I _{el} ⁽³⁾ mBq·m ⁻³	¹³¹ I _{or} ⁽⁴⁾ mBq·m ⁻³	¹³⁷ Cs mBq·m ⁻³	Nat. ⁽⁵⁾ mBq·m ⁻³
03/02/16	< 0.046	< 0.2	< 0.6	< 0.035	< 1.6
02/03/16	< 0.053	< 0.2	< 0.6	< 0.037	< 1.5
06/04/16	< 0.039	< 0.1	< 0.4	< 0.029	< 1.5
03/05/16	< 0.050	< 0.2	< 0.7	< 0.038	< 1.8
07/06/16	< 0.041	< 0.2	< 0.6	< 0.031	< 1.6
06/07/16	< 0.042	< 0.2	< 0.5	< 0.031	< 1.3
03/08/16	< 0.049	< 0.2	< 0.8	< 0.038	< 1.8
07/09/16	< 0.052	< 0.1	< 0.7	< 0.039	< 1.9
06/10/16	< 0.052	< 0.3	< 0.7	< 0.041	< 2
03/11/16	< 0.08	< 0.3	< 1	< 0.058	< 3
05/12/16	< 0.046	< 0.1	< 0.4	< 0.054	< 2
05/01/17	< 0.050	< 0.1	< 0.4	< 0.034	< 2

⁽¹⁾ Analysis was performed on a combined sample of the monthly samples taken from all five locations (21, 22, 23, 27 and 29).

⁽²⁾ End date of monthly sampling period.

⁽³⁾ Elemental ¹³¹I.

⁽⁴⁾ Organically bound ¹³¹I.

⁽⁵⁾ Naturally occurring γ -emitters.

Table A18 Activity concentrations of γ -emitters in grass near the Borssele nuclear power plant in 2016 ⁽¹⁾

Date	Mass kg·m ⁻²	⁶⁰ Co Bq·kg ⁻¹ (2)	¹³¹ I Bq·kg ⁻¹ (2)	¹³⁷ Cs Bq·kg ⁻¹ (2)
03/02/16	0.664	< 2	< 2	< 1
02/03/16	0.622	< 2	< 2	< 2
06/04/16	0.550	< 1	< 0.9	< 0.9
03/05/16	0.711	< 2	< 2	< 1
07/06/16	0.880	< 2	< 1	< 1
06/07/16	0.733	< 2	< 1	< 1
03/08/16	0.835	< 2	< 2	< 2
07/09/16	1.21	< 2	< 2	< 2
06/10/16	1.15	< 1	< 0.9	< 0.8
03/11/16	0.703	< 3	< 2	< 2
05/12/16	1.28	< 0.8	< 1	< 0.7
05/01/17	0.604	< 2	< 1	< 1

⁽¹⁾ Analysis was performed on a combined sample of the monthly samples taken from all five locations (21, 22, 23, 27 and 29).

⁽²⁾ Dry weight.

Table A19 Activity concentrations of γ -emitters in soil near the Borssele nuclear power plant in 2016 ⁽¹⁾

Location	Mass kg·m ⁻²	⁵⁴ Mn Bq·kg ⁻¹ (2)	⁶⁰ Co Bq·kg ⁻¹ (2)	¹³⁴ Cs Bq·kg ⁻¹ (2)	¹³⁷ Cs Bq·kg ⁻¹ (2)
O1	76.8	< 0.3	< 0.4	< 0.2	1.27 ± 0.06
O2	91.2	< 0.4	< 0.3	< 0.3	1.48 ± 0.07
O3	73.8	< 0.3	< 0.3	< 0.4	1.22 ± 0.06
O4	71.2	< 0.3	< 0.3	< 0.3	1.47 ± 0.06

⁽¹⁾ Analysis was performed on four samples taken near the outlet of the plant on 9 May 2016.

⁽²⁾ Dry weight.

Table A20 Residual β activity concentrations in water from the Western Scheldt in 2016

Date	Residual β Bq·L ⁻¹			
	1	2	3	4
03/02/16	0.043 ± 0.008	0.034 ± 0.007	0.018 ± 0.009	0.052 ± 0.007
02/03/16	0.060 ± 0.008	0.037 ± 0.007	0.020 ± 0.008	0.068 ± 0.006
06/04/16	0.050 ± 0.007	0.035 ± 0.007	0.040 ± 0.006	0.066 ± 0.007
03/05/16	0.033 ± 0.008	0.027 ± 0.007	0.011 ± 0.013	0.033 ± 0.010
07/06/16	0.035 ± 0.008	0.051 ± 0.007	0.068 ± 0.010	0.029 ± 0.007
06/07/16	0.035 ± 0.007	0.050 ± 0.006	0.033 ± 0.007	0.052 ± 0.007
03/08/16	0.037 ± 0.008	0.023 ± 0.007	0.024 ± 0.005	0.061 ± 0.008
07/09/16	0.030 ± 0.009	0.013 ± 0.008	0.049 ± 0.008	0.018 ± 0.007
06/10/16	0.049 ± 0.009	0.030 ± 0.008	0.060 ± 0.009	0.033 ± 0.008
03/11/16	0.036 ± 0.008	0.132 ± 0.010	0.028 ± 0.008	0.052 ± 0.008
05/12/16	0.060 ± 0.009	0.061 ± 0.010	0.116 ± 0.009	0.060 ± 0.010
05/01/17	0.085 ± 0.012	0.054 ± 0.008	< 0.021	0.036 ± 0.008

Table A21 ^3H activity concentrations in water from the Western Scheldt in 2016

Date	^3H Bq·L ⁻¹			
Location	1	2	3	4
03/02/16	5.6 ± 1.2	5.1 ± 1.1	2.7 ± 1.1	7.0 ± 1.2
02/03/16	4.5 ± 1.1	5.0 ± 1.1	4.6 ± 1.1	6.1 ± 1.2
06/04/16	7.0 ± 1.2	5.6 ± 1.2	6.3 ± 1.2	5.6 ± 1.2
03/05/16	5.7 ± 1.1	6.9 ± 1.2	3.9 ± 1.1	3.5 ± 1.1
07/06/16	1.8 ± 1.1	2.7 ± 1.1	4.3 ± 1.1	3.3 ± 1.1
06/07/16	2.6 ± 1.1	4.7 ± 1.2	1.4 ± 1.1	2.5 ± 1.1
03/08/16	5.6 ± 1.1	5.2 ± 1.1	3.0 ± 1.1	5.5 ± 1.1
07/09/16	4.2 ± 1.2	2.9 ± 1.1	3.9 ± 1.2	2.7 ± 1.1
06/10/16	5.7 ± 1.2	3.7 ± 1.1	3.6 ± 1.2	6.7 ± 1.2
03/11/16	3.3 ± 1.1	4.6 ± 1.1	4.7 ± 1.1	3.6 ± 1.1
05/12/16	4.4 ± 1.1	7.3 ± 1.2	4.2 ± 1.1	4.3 ± 1.1
05/01/17	5.1 ± 1.1	4.9 ± 1.1	2.3 ± 0.8	5.8 ± 1.2

Table A22 Gross β activity concentrations in suspended solids from the Western Scheldt in 2016

Date	Gross β - kBq·kg ⁻¹			
Location	1	2	3	4
03/02/16	1.8 ± 0.9	2.2 ± 0.4	1.13 ± 0.09	3.1 ± 0.9
02/03/16	1.6 ± 0.4	2.0 ± 0.3	1.6 ± 0.4	1.8 ± 0.3
06/04/16	1.5 ± 0.8	2.3 ± 0.5	1.50 ± 0.14	2.1 ± 0.3
03/05/16	2.3 ± 0.7	2.1 ± 0.7	1.0 ± 0.5	0.2 ± 1.2
07/06/16	1.51 ± 0.10	1.50 ± 0.13	2.6 ± 0.5	1.76 ± 0.16
06/07/16	2.4 ± 0.5	3.0 ± 0.6	2.7 ± 0.9	2.6 ± 0.5
03/08/16	1.0 ± 0.3	2.3 ± 0.5	1.2 ± 0.3	2.7 ± 0.6
07/09/16	1.4 ± 0.8	2.3 ± 0.9	2.4 ± 0.8	1.8 ± 0.6
06/10/16	1.2 ± 0.3	1.5 ± 0.2	0.7 ± 0.3	1.1 ± 0.2
03/11/16	2.1 ± 0.5	1.7 ± 0.6	2.0 ± 0.3	1.4 ± 0.2
05/12/16	1.7 ± 0.4	0.63 ± 0.19	0.3 ± 0.2	0.6 ± 0.4
05/01/17	1.3 ± 1.2	2.6 ± 1.2	1.2 ± 1.3	2.7 ± 1.0

Table A23 Activity concentrations of γ -emitters in seaweed from the Western Scheldt in 2016 ⁽¹⁾

Date	Mass kg	⁶⁰ Co Bq·kg ⁻¹ (2)	¹³¹ I Bq·kg ⁻¹ (2)	¹³⁷ Cs Bq·kg ⁻¹ (2)
03/02/16	0.253	< 1	< 0.8	< 0.7
02/03/16	0.167	< 1	< 1	< 1
06/04/16	0.236	< 0.6	< 0.5	< 0.5
03/05/16	0.115	< 2	< 2	< 1
07/06/16	0.111	< 1	< 2	< 1
06/07/16	0.119	< 1	< 2	< 1.0
03/08/16	0.102	< 2	< 1	< 2
07/09/16	0.142	< 1	< 1.0	< 0.7
06/10/16	0.15	< 0.7	< 0.8	< 0.8
03/11/16	0.096	< 2	< 2	< 2
05/12/16	0.166	< 1	< 1	< 0.9
05/01/17	0.117	< 2	< 2	< 1

⁽¹⁾ Analysis was performed on a combined sample of the monthly samples taken from all four locations (1, 2, 3 and 4).

⁽²⁾ Dry weight.

Table A24 Activity concentrations of γ -emitters in sediment from the Western Scheldt in 2016 ⁽¹⁾

Location	Mass kg·m ⁻²	⁶⁰ Co Bq·kg ⁻¹ (2)	¹³¹ I Bq·kg ⁻¹ (2)	¹³⁷ Cs Bq·kg ⁻¹ (2)
03/02/16	67.2	< 0.2	< 0.3	0.36 ± 0.03
02/03/16	79.9	< 0.2	< 0.2	0.27 ± 0.03
06/04/16	77.5	< 0.2	< 0.2	0.44 ± 0.07
03/05/16	74.4	< 0.2	< 0.2	0.29 ± 0.03
07/06/16	63.5	< 0.3	< 0.2	0.77 ± 0.05
06/07/16	76.7	< 0.2	< 0.3	0.39 ± 0.04
03/08/16	78.9	< 0.2	< 0.2	0.38 ± 0.03
07/09/16	59.7	< 0.2	< 0.3	0.99 ± 0.05
06/10/16	57.6	< 0.2	< 0.3	< 0.2
03/11/16	65.2	< 0.3	< 0.3	0.81 ± 0.05
05/12/16	69.8	< 0.2	< 0.2	0.25 ± 0.03
05/01/17	61.8	< 0.2	< 0.3	0.84 ± 0.04

⁽¹⁾ Analysis was performed on a combined sample of the monthly samples from all four locations (1, 2, 3 and 4).

⁽²⁾ Dry weight.

Appendix B Presentation of data

The methods described below were applied to the data provided by the RIVM/VLH (e.g. air dust and deposition), unless otherwise stated. Data from the other institutions are reported as provided, unless otherwise stated.

B.1 Correction for radioactive decay

In general, the activities of specific radionuclides are corrected for radioactive decay. The activities measured in the sample are multiplied by a decay factor, combining the time from halfway through the sampling period to the time of analysis, the decay during the measurement and the half-life of the radionuclide. If the radionuclides are unknown, as with gross α and gross β , no correction for radioactive decay is made.

B.2 Calculation of sums and averages

In the calculation of weekly, monthly or yearly averages or sums, the original results before rounding-off are used. If a certain radionuclide cannot be measured, the detection limit is used in the calculation of the sums. In that case, only a range (lower and upper limit) is given instead of a total with an uncertainty. Both range and total with an uncertainty are presented with a 68% confidence interval.

The lower and upper limits are calculated as follows:

$$\text{Lower limit} = \sum x_i - \sqrt{\sum s_i^2}$$

$$\text{Upper limit} = \sum x_i + \sqrt{\sum s_i^2} + \sum \text{MDA}_j$$

where

x_i = weekly or monthly result that is not a detection limit;

$\sqrt{\sum s_i^2}$ = the uncertainty in the sum;

s_i = uncertainty in the weekly or monthly result (1σ);

MDA_j = weekly or monthly result that is a detection limit.

The detection limits are omitted in the calculation of the averages. If data are not reported (e.g. a sample was not analysed/available), the sampling period is not taken into account for the calculation of the sum or average.

B.3 Calculation of uncertainties

The uncertainties given in Tables A1–A8 are a combination of the statistical uncertainties and estimations of the experimental uncertainties. In the yearly total, the uncertainty is the square root of the sum of the squared weekly or monthly uncertainties. In the yearly

average, the uncertainty is the square root of the sum of the squared weekly uncertainties divided by the number of weeks.

B.4 Mixed diets: conversion from Bq·kg⁻¹ to intake in Bq·day⁻¹

With respect to the results presented for mixed diets (Chapter 8), RIKILT Wageningen UR used food consumption patterns to convert the measured concentrations of ⁹⁰Sr, ¹³⁴Cs and ¹³⁷Cs in food (Bq·kg⁻¹) to an average daily intake value per person per day (Bq·day⁻¹). For the Netherlands, the food consumption patterns were investigated by the RIVM and the results can be found in the report 'Dutch National Food Consumption Survey 2007–2010. Diet of children and adults aged 7 to 69 years' [76]. In this report, the consumption patterns are presented by food category, sex and age group, in grams per consumption day, and the percentage of consumption days. For the calculations in the current report, these values were combined to produce an average consumption amount in g·day⁻¹ for each food category, sex and age group.

For each sex and age group and specific radionuclide, the daily intake ($DI_{a,s,n}$) is then calculated as follows:

$$DI_{a,s,n} = \sum_i \frac{DI_{a,s,i}}{1000} \times AC_{i,n}$$

where

$DI_{a,s,n}$ = daily intake per age group, sex and radionuclide (Bq·day⁻¹);

i = food category;

$DI_{a,s,i}$ = daily intake per age group, sex and food category (g·day⁻¹);

$AC_{i,n}$ = activity concentration per food category and radionuclide (Bq·kg⁻¹).

These daily intakes were then averaged over the different age groups and sexes to obtain the total daily intake per person for each radionuclide. To include the monitoring results of ready meals in the final result, the assumption was made that ready meals make up 10% of the consumption of the categories meat, fish and vegetables.

Appendix C Glossary

Ambient dose equivalent	An operational quantity used when monitoring radiation in the environment. The unit of ambient dose equivalent is the Sievert (Sv).
Becquerel (Bq)	One radioactive transformation per second.
Decay product	A decay product (also known as a daughter product, daughter isotope, or daughter nuclide) is a nuclide resulting from the radioactive decay of a parent isotope or precursor nuclide. The decay product may be stable or it may decay to form a daughter product of its own.
Dose rate	The radiation dose delivered per unit of time.
Effective dose	The sum of the equivalent doses from internal and external radiation in all tissue and organs of the body, having been weighted by their tissue weighting factors. The unit of effective dose is the Sievert (Sv).
Gross alpha activity	Gross α (or total α) activity is the total activity of radionuclides emitting α radiation.
Gross beta activity	Gross β (or total β) activity is the total activity of radionuclides emitting β radiation. Depending on the measurement methodology, it might exclude tritium and/or radon daughters.
Limit	Consumption limits in milk, foodstuffs and of feeding stuffs set in European legislation [65, 66, 67].
Parametric value	The value of radioactive substances in drinking water above which Member States shall assess whether the presence of radioactive substances in water intended for human consumption poses a risk to human health that requires action and, where necessary, shall take remedial action to improve the quality of water to a level which complies with the requirements for the protection of human health from a radiation protection point of view [48].
Radioactivity	The emission of α particles, β particles, neutrons and γ or X radiation from the disintegration of an atomic nucleus. The unit of radioactivity is the Becquerel (Bq).
Radiocesium activity	Sum of the activity of ^{134}Cs and ^{137}Cs .
Radionuclide	An unstable form of an element that undergoes radioactive decay.

Residual beta activity	The residual β activity is the gross β activity (total β activity) minus the β activity of naturally occurring ^{40}K . For brackish and salt water, the RWS uses a direct method to determine residual β activity [43].
Screening level	Level for gross α or gross β activity in drinking water ($\text{Bq}\cdot\text{L}^{-1}$) below which it can be assumed that the individual dose is less than the parametric value of 0.1 mSv and radiological investigation is not needed unless it is known from other sources of information that specific radionuclides are present in the water that are liable to cause an individual dose in excess of 0.1 mSv [48].

References

- [1] EC, 2000. Recommendation of the Commission of the European Communities on the Application of Article 36 of the Euratom Treaty. EC Brussels, 2000/473/Euratom.
- [2] R.B. Tax, P.J.M. Kwakman, A.P.P.A. van Lunenburg, M.H. Tijsmans, 1994. Development of a High Volume Air Sampler for the sensitive detection of γ -emitting radionuclides attached to aerosols. Results obtained in the test period 1991–1992. RIVM Bilthoven, Report no. 610056005.
- [3] G.J. Knetsch, 2011. Afnametest Snow White. RIVM Bilthoven, internal report.
- [4] NEN, 2009. NEN 5621, Radioactivity measurements – Sampling of airborne substances. NEN, Delft.
- [5] NEN, 2006. NEN 5636, Radioactivity measurements – Determination of artificial gross-alpha activity, artificial gross-beta activity and gamma spectrometry of air filters and calculation of the volumic activity of sampled air. NEN, Delft.
- [6] NEN, 2002. NEN 5623, Radioactivity measurements – Determination of the activity of gamma ray-emitting nuclides in a counting sample by semiconductor gamma spectrometry. NEN, Delft.
- [7] G.J. Knetsch (ed.), 2006. Environmental radioactivity in the Netherlands. Results in 2005. RIVM Bilthoven, Report no. 861020013.
- [8] S. Sugihara, N. Momoshima, Y. Maeda, S. Osaki, 2000. Variation of atmospheric ^7Be and ^{210}Pb depositions at Fukuoka, Japan. IRPA 10th congress, Web page: www2000.irpa.net/irpa10/cdrom/00822.pdf (January 2019).
- [9] C. Ródenas, J. Gómez, L.S. Quindós, P.L. Fernández, J. Soto, 1997. ^7Be concentrations in air, rainwater and soil in Cantabria (Spain). *Applied Radiation and Isotopes* 48, 545–548.
- [10] S. Talpos and V. Cuculeanu, 1997. A study of the vertical diffusion of ^7Be in the atmosphere. *Journal of Environmental Radioactivity* 36 (1), 93–106.
- [11] K.N. Yu and L.Y.L. Lee, 2002. Measurements of atmospheric ^7Be properties using high-efficiency gamma spectroscopy. *Applied Radiation and Isotopes* 57, 941–946.
- [12] C. Papastefanou and A. Ioannidou, 1995. Aerodynamic size association of ^7Be in ambient aerosols. *Journal of Environmental Radioactivity* 26, 273–282.
- [13] H.W. Feely, R.J. Larsen, C.G. Sanderson, 1989. Factors that cause seasonal variations in ^7Be concentrations in surface air. *Journal of Environmental Radioactivity* 9, 223–249.
- [14] C.L. Fogh, J. Roed, K.G. Andersson, 1999. Radionuclide resuspension and mixed deposition at different heights. *Journal of Environmental Radioactivity* 46, 67–75.
- [15] Solar Cycle Progression. Web page: www.swpc.noaa.gov/products/solar-cycle-progression (January 2019).

- [16] R.C.G.M. Smetsers and R.O. Blaauboer, 1997. A dynamic compensation method for natural ambient dose rate based on 6 years' data from the Dutch Radioactivity Monitoring Network. *Radiation Protection Dosimetry* 69 (1), 19–31.
- [17] S.I. Dusha-Gudym, 2005. Transport of radioactive materials by wildland fires in the Chernobyl accident zone: how to address the problem. *International Forest Fire News* 32. Web page: gfmcc.org/online/wp-content/uploads/20-Dusha-Gudym-3.pdf (January 2019).
- [18] KNMI database. Web page: www.knmi.nl/klimatologie (January 2019).
- [19] T. Hantke, F.J. Aldenkamp, R.M.W. Overwater, H. Slaper, 1998. De jacht op een ^{137}Cs -wolk in Zuid-Europa – 'aftermath' van een ongeval in Algeciras. *NVS Nieuws* 23 (4).
- [20] G.J. Knetsch (ed.), 2013. Environmental radioactivity in the Netherlands. Results in 2011. RIVM Bilthoven, Report no. 610891004.
- [21] UNSCEAR, 2000. Sources and effects of ionizing radiation. Volume 1: Sources.
- [22] E.A. Bondietti, C. Papastefanou, C. Rangarajan, 1987. Aerodynamic size associations of natural radioactivity with ambient aerosols. In: *Radon and its Decay Products: Occurrence, Properties and Health Effects*, ACS Symposium Series No. 331, P.K. Hopke (ed.), American Chemical Society, Washington, DC, 377–397.
- [23] T. Suzuki, Y. Maruyama, N. Nakayama, K. Yamada, K. Ohta, 1999. Measurement of $^{210}\text{Po}/^{210}\text{Pb}$ activity ratio in size fractionated aerosols from the coast of the Japan Sea. *Atmospheric Environment* 33, 2285–2288.
- [24] T. Tokieda, K. Yamanaka, K. Harada, S. Tsunogai, 1996. Seasonal variations of residence time and upper atmospheric contribution of aerosols studied with Pb-210, Bi-210, Po-210 and Be-7. *Tellus*, 48B, 690–702.
- [25] G. Lambert, P. Bristeau, G. Polian, 1976. Emission and enrichments of radon daughters from Etna volcano magma. *Geophysical Research Letters* 3 (12), 724–726.
- [26] J. Sato, T. Doi, T. Segawa, S. Sugawara, 1994. Seasonal variation of atmospheric concentrations of ^{210}Pb and ^7Be at Tsukuba, Japan, with a possible observation of ^{210}Pb originating from the 1991 eruption of the Pinatubo volcano, Philippines. *Geochemical Journal* 28, 123–129.
- [27] J.P. Beks, D. Eisma, J. van der Plicht, 1998. A record of atmospheric ^{210}Pb deposition in the Netherlands. *Science of the Total Environment* 222, 35–44.
- [28] J. Sato, 2003. Natural radionuclides in volcanic activity. *Applied Radiation and Isotopes* 58, 393–399.
- [29] G. Lambert, B. Ardouin, G. Polian, 1982. Volcanic output of long-lived radon daughters. *Journal of Geophysical Research* 87 (C13), 11103–11108.

- [30] C.P. Tanzi and G.J. Knetsch, 2012. ^{210}Po and ^{210}Pb in the Netherlands: releases to air from industrial plants compared to environmental monitoring. Proceedings of EU-NORM 1st International Symposium, 5–8 June, Tallinn, Estonia, 59–66.
- [31] E.Y. Nho, B. Ardouin, M.F. Le Cloarec, M. Ramonet, 1996. Origins of ^{210}Po in the atmosphere at Lamto, Ivory Coast: biomass burning and Saharan dust. *Atmospheric Environment* 30 (22), 3705–3714.
- [32] M.A. Mélières, M. Pourchet, S. Richard, 2003. Surface air concentration and deposition of lead-210 in French Guiana: two years of continuous monitoring. *Journal of Environmental Radioactivity* 66, 261–269.
- [33] P.G. Appleby, A.O. Koulikov, L. Camarero, M. Ventura, 2002. The input and transmission of fall-out radionuclides through Redó, a high mountain lake in the Spanish Pyrenees. *Water, Air & Soil Pollution: Focus* 2, 19–31.
- [34] NEN, 2009. NEN 5628, Radioactivity measurements – Sampling of the Deposition. NEN, Delft.
- [35] NEN, 2006. NEN 5622, Radioactivity measurements – Determination of massic gross-alpha activity of a solid counting sample by the thick source method. NEN, Delft.
- [36] NEN, 2016. NEN 5627, Radioactivity measurements – Determination of massic gross beta activity and massic rest beta activity of a solid counting sample. NEN, Delft.
- [37] NEN, 2007. NEN 5694, Radioactivity measurements – Methods for radiochemical determination of polonium-210 and lead-210. NEN, Delft.
- [38] C.J.W. Twenhöfel, C. de Hoog van Beynen, A.P.P.A. van Lunenburg, G.J.E. Slagt, R.B. Tax, P.J.M. van Westerlaak, F.J. Aldenkamp, 2005. Operation of the Dutch 3rd Generation National Radioactivity Monitoring Network. In: *Automatic Mapping Algorithms for Routine and Emergency Monitoring Data, Spatial Interpolation Comparison 2004* by IES, G. Dubois (ed.), European Committee, JRC, EUR 21595 2005, 19–31.
- [39] R.O. Blaauboer and R.C.G.M. Smetsers, 1996. Variations in outdoor radiation levels in the Netherlands. Thesis University of Groningen, Groningen.
- [40] C. de Hoog and R.B. Tax, 2003. Achtergronddocument bij NMR integrale rapportage 2002. RIVM Bilthoven, internal report.
- [41] Federal Aviation Administration. Web page: www.faa.gov/data_research/research/med_humanfacs/aeromedical/radiobiology/heliocentric/ (last accessed on 21 January 2019).
- [42] E.J. de Jong and O.C. Swertz, 2000. Radioactieve stoffen in de zoute wateren. RIKZ, The Hague, Report no. RIKZ/2000.041.
- [43] L.J. Gilde, K.H. Prins, C.A.M. van Helmond, 1999. Monitoring zoete rijkswateren. RIZA Lelystad, Report no. 99.004.
- [44] M.W.M. Bogaart-Scholte et al., 2014. MWTL Meetplan 2014 – Monitoring waterstaatkundige toestand des lands milieumeetnet rijkswateren chemie en biologie. Rijkswaterstaat Waterdienst, Lelystad.

- [45] Rijkswaterstaat. Web page: http://live.waterbase.nl/waterbase_wns.cfm?taal=nl (August 2018).
- [46] H. Schuijn, 2009. Analyseboek monitoring zoet, versie 1 juni 2009. Rijkswaterstaat Waterdienst, Lelystad.
- [47] H. Schuijn, 2009. Analyseboek monitoring zout, versie 1 juni 2009. Rijkswaterstaat Waterdienst, Lelystad.
- [48] EC, 2013. Council Directive laying down requirements for the protection of the health of the general public with regard to radioactive substances in water intended for human consumption. EC Brussels, 2013/51/EURATOM.
- [49] H.A.J.M. Reinen, C. de Hoog, F. Wetsteyn, J.G.M.M. Smeenk, H.A.M. Ketelaars, A.D. Hulsmann, J.M. van Steenwijk, A.J. Stortenbeek, 2003. Meetstrategie drinkwater bij kernongevallen. VROM-Inspectie Den Haag, Report No. 15060/177.
- [50] Dutch Drinking Water Decree (2001); Staatsblad nr 31, until 30 June 2011.
- [51] Dutch Drinking Water Decree (2011), from 1 July 2011 onwards.
- [52] G.J. Knetsch (ed.), 2003. Monitoring of radiation in the environment in the Netherlands. Results in 2002. RIVM Bilthoven, Report no. 861020005.
- [53] G.J. Knetsch (ed.), 2004. Monitoring of radiation in the environment in the Netherlands. Results in 2003. RIVM Bilthoven, Report no. 861020010.
- [54] G.J. Knetsch (ed.), 2005. Monitoring of radiation in the environment in the Netherlands. Results in 2004. RIVM Bilthoven, Report no. 861020011.
- [55] G.J. Knetsch (ed.), 2007. Environmental radioactivity in the Netherlands. Results in 2006. RIVM Bilthoven, Report no. 610791001.
- [56] G.J. Knetsch (ed.), 2008. Environmental radioactivity in the Netherlands. Results in 2007. RIVM Bilthoven, Report no. 610791002.
- [57] G.J. Knetsch (ed.), 2010. Environmental radioactivity in the Netherlands. Results in 2008. RIVM Bilthoven, Report no. 610791003.
- [58] M.C.E. Groot and G.J. Knetsch (eds), 2011. Environmental radioactivity in the Netherlands. Results in 2009. RIVM Bilthoven, Report no. 610891002.
- [59] G.J. Knetsch (ed.), 2012. Environmental radioactivity in the Netherlands. Results in 2010. RIVM Bilthoven, Report no. 610891003.
- [60] G.J. Knetsch (ed.), 2014. Environmental radioactivity in the Netherlands. Results in 2012. RIVM Bilthoven, Report no. 610891005.
- [61] G.J. Knetsch (ed.), 2015. Environmental radioactivity in the Netherlands. Results in 2013. RIVM Bilthoven, Report no. 2015-0040.
- [62] G.J. Knetsch (ed.), 2016. Environmental radioactivity in the Netherlands. Results in 2014. RIVM Bilthoven, Report no. 2016-0182.
- [63] G.J. Knetsch (ed.), 2017. Environmental radioactivity in the Netherlands. Results in 2015. RIVM Bilthoven, Report no. 2016-0183.

- [64] P.J.M. Kwakman, J.F.M. Versteegh, 2015. Radon-222 in groundwater and finished drinking water in the Dutch provinces Overijssel and Limburg. Measuring campaign 2015. RIVM Bilthoven, RIVM Letter report no. 2016-0048.
- [65] EC, 2009. Council Regulation on the conditions governing imports of agricultural products originating in third countries following the accident at the Chernobyl nuclear power station. EC Brussels, No. 1048/2009.
- [66] EC, 2008. Council Regulation on the conditions governing imports of agricultural products originating in third countries following the accident at the Chernobyl nuclear power station. EC Brussels, No. 733/2008.
- [67] EC, 1989. Council Regulation amending Regulation (Euratom) No. 3954/87 laying down maximum permitted levels of radioactive contamination of foodstuffs and of feeding stuffs following a nuclear accident or any other case of radiological emergency. EC Brussels, No. 2218/89.
- [68] EC, 2013. Council Directive laying down basic safety standards for protection against the dangers arising from exposure to ionising radiation, and repealing Directives 89/618/Euratom, 90/641/Euratom, 96/29/Euratom, 97/43/Euratom and 2003/122/Euratom. EC Brussels, No. 2013/59/Euratom.
- [69] Keuringsdienst van Waren Oost, 1998. Werkvoorschrift CHE01-OT802, Keuringsdienst van Waren Oost, Bepaling van de activiteit van gammastraling uitzendende nucliden in een telmonster met halfgeleiderspectrometrie. Nijmegen.
- [70] Keuringsdienst van Waren Oost, 2003. Werkvoorschrift CHE01-WV143, Keuringsdienst van Waren Oost, Bepaling van de activiteit van gammastraling uitzendende nucliden in een telmonster met de LMRV voedselmonitor. Zutphen.
- [71] Jaarverslag NPK 2016. Nederlandse Voedsel en Waren Autoriteit.
- [72] J.J. Donk, 2016. Resultaten van de dosistempo- en radioactiviteitsmetingen in de omgeving van Borssele over het jaar 2016. NRG Arnhem, Report No. 23868/17.142118 (commissioned by N.V. EPZ).
- [73] KEMA, 1994. Uitgangspunten voor de omgevingsbewakingsprogramma's van de kerncentrales te Dodewaard en Borssele. KEMA Arnhem, Report no. 40318/40575-NUC 94-5935.
- [74] J.E.M. Jacobs (ed.), 2001. Monitoring of radiation in the Environment. Results in the Netherlands in 1999. RIVM Bilthoven, Report no. 610056046.
- [75] Letter to the State Health Inspectorate of the Ministry of Housing, Spatial Planning and Environment, 13 January 1997. Reference no. 23/97 LSO Le/Ald/jdk. Onderzoek naar de slechte resultaten in 1995 van de bepaling van ^{210}Po en ^{210}Pb in natte en droge depositie. RIVM Bilthoven, internal report.
- [76] C.T.M. van Rossum, H.P. Fransen, J. Verkaik-Kloosterman, E.J.M. Buurma-Rethans, M.C. Ocke, 2011. Dutch National Food Consumption Survey 2007–2010: Diet of children and adults aged 7 to 69 years. RIVM Bilthoven, Report no. 350050006.

Erratum

Author C.P. Tanzi

Date 13 June 2019

Report number 2018-0160

Report title

Environmental radioactivity in the Netherlands - Results in 2016

Errors

Two transcription errors have been identified in the Summary Table S.1 on page 15: for drinking water, the number of locations for ^3H is 180 (instead of the reported 88), and for ^{90}Sr in milk, the value is $<0.2 \text{ Bq}\cdot\text{kg}^{-1}$ (instead of the reported $<5 \text{ Bq}\cdot\text{kg}^{-1}$). The correct table is given below.

There is typographical error in the caption of table A10 on page 94, where the year 2017 is incorrectly mentioned, instead of the relevant year 2016.

The correct caption is given below:

Table A10 Yearly average ambient dose equivalent rate for the NMR stations in 2016

There is typographical error in note (2) of table A10 on page 95, where the year 2018 is incorrectly mentioned, instead of the relevant year 2016. The correct note is given below:

(2) A few stations have been moved during 2016: Wekerom station was moved to Otterlo, Harlingen station to Witmarsum, Hengelo (Ov) station to Laren (Gld) and Den Ham station to Vroomshoop.

There is typographical error in the caption of table A12 on page 99, where the year 2015 is incorrectly mentioned, instead of the relevant year 2016.

The correct caption is given below:

Table A12 ^{60}Co , ^{131}I , ^{137}Cs and ^{210}Pb activity concentrations in suspended solids ($\text{Bq}\cdot\text{kg}^{-1}$) in surface water in 2016 as measured by RWS

These corrections have no consequence for the conclusions of the report.

Table S1 Summary of the results from the Dutch monitoring programme in 2016

Matrix	Parameter	Locations	Values	Frequency (per year)
Air dust ⁽¹⁾	Gross α	1	0.019 mBq·m ⁻³	52
	Gross β	1	0.362 mBq·m ⁻³	52
	⁷ Be	1	2.910 mBq·m ⁻³	52
	¹³⁷ Cs	1	0.000208 mBq·m ⁻³	52
	²¹⁰ Pb	1	0.312 mBq·m ⁻³	52
Deposition ⁽²⁾	Gross α	1	66 Bq·m ⁻²	12
	Gross β	1	88 Bq·m ⁻²	12
	³ H ⁽³⁾	1	22–1,510 Bq·m ⁻²	12
	⁷ Be	1	1,375 Bq·m ⁻²	52
	¹³⁷ Cs ⁽³⁾	1	0.04–1.15 Bq·m ⁻²	52
	²¹⁰ Pb	1	95–98 Bq·m ⁻²	52
	²¹⁰ Po	1	40.6 Bq·m ⁻²	12
Surface water ⁽¹⁾	Gross α	8	29–249 mBq·L ⁻¹	12–13 ⁽⁴⁾
	Residual β	8	13–112 mBq·L ⁻¹	12–13 ⁽⁴⁾
	³ H	8	1,550–14,600 mBq·L ⁻¹	6–13 ⁽⁴⁾
	⁹⁰ Sr	3	< 1–3.1 mBq·L ⁻¹	6–7 ⁽⁴⁾
	²²⁶ Ra	4	4.0–33.1 mBq·L ⁻¹	6–7 ⁽⁴⁾
Suspended solids in surface water ⁽¹⁾	⁶⁰ Co	8	< 1–10.4 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	¹³¹ I	8	< 1–36.0 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	¹³⁷ Cs	8	3.0–12.4 Bq·kg ⁻¹	4–51 ⁽⁴⁾
	²¹⁰ Pb	4	80–140 Bq·kg ⁻¹	6–7 ⁽⁴⁾
Seawater ⁽¹⁾	Gross α	8	190–420 mBq·L ⁻¹	4–13 ⁽⁴⁾
	Residual β	8	34–140 mBq·L ⁻¹	4–13 ⁽⁴⁾
	³ H	8	350–5,200 mBq·L ⁻¹	4–13 ⁽⁴⁾
	⁹⁰ Sr	4	< 1.4–1.9 mBq·L ⁻¹	4–13 ⁽⁴⁾
Suspended solids in seawater ⁽¹⁾	¹³⁷ Cs	1	3.3 Bq·kg ⁻¹	4 ⁽⁴⁾
	²¹⁰ Pb	1	62 Bq·kg ⁻¹	4 ⁽⁴⁾
Drinking water ⁽¹⁾	Gross α	183	< 0.06 Bq·L ⁻¹	388 ⁽⁵⁾
	Gross β	186	< 0.1 Bq·L ⁻¹	428 ⁽⁵⁾
	Residual β	168	< 0.1 Bq·L ⁻¹	390 ⁽⁵⁾
	³ H	180	< 4.0 Bq·L ⁻¹	448 ⁽⁵⁾
Milk ⁽¹⁾	⁴⁰ K	22	51.2 Bq·kg ⁻¹ ⁽⁶⁾	581 ⁽⁵⁾
			50.1 Bq·kg ⁻¹ ⁽⁷⁾	11
	⁶⁰ Co	22	< 1.4 Bq·kg ⁻¹	581 ⁽⁵⁾
	⁹⁰ Sr	22	< 0.2 Bq·kg ⁻¹	47 ⁽⁵⁾
	¹³¹ I	22	< 0.6 Bq·kg ⁻¹	581 ⁽⁵⁾
	¹³⁴ Cs	22	< 0.6 Bq·kg ⁻¹	581 ⁽⁵⁾
	¹³⁷ Cs	22	< 0.5 Bq·kg ⁻¹	581 ⁽⁵⁾

Paraaf projectleider/labhoofd

Voor akkoord, 28 juni 2019

GJ Knetsch

Programma coördinator "Uitvoering Wettelijke Taken"

.....

C.P. Tanzi

.....

RIVM Report 2018-0160

Published by:

**National Institute for Public Health
and the Environment, RIVM**
P.O. Box 1 | 3720 BA Bilthoven
The Netherlands
www.rivm.nl/en

Januari 2019

Committed to *health and sustainability*