


Vaccine Coverage Dutch National Influenza Prevention Program 2019: Brief monitor

M.J. Heins, M. Hooiveld, J.C. Korevaar

Further increase in vaccine coverage in 2019

For the second year in row there was a small increase in influenza vaccine coverage. During the yearly influenza vaccination campaign of 2019, 52.6% of the target population was vaccinated. In 2017 this was 49.9%. The target population are persons with an increased risk for complications due to influenza because of age or a medical condition. The highest vaccine coverage (70.6%) was found in persons over 65 years of age who also had a medical indication for vaccination. Vaccine coverage was lowest in persons between 60 and 64 years old without a medical indication for vaccination (29.5%). Vaccine coverage was highest in persons with cardiovascular diseases (63.4%) and lowest among persons with HIV (44.9%).

Figure 1 Vaccine coverage by age group and medical indication, 1996-2019



* In 2008, the age group 60 – 64 years was included among the target groups for influenza vaccination. Therefore, the vaccination coverage of the entire target population from 2008 onwards cannot be compared to previous years.

Results from 1996-2013: Tacken M.A., et al. Monitoring vaccinatiegraad Nationaal Programma Grieppreventie 2013. Nijmegen, IQ Healthcare: 2014.

Table 1 Size and influenza vaccine coverage by age group and medical indication (2019)

Age group	Size (% of population)	Vaccine coverage (% of population)
0-59 years with medical indication*	6.7	32.0
60 years and older	26.4	54.8
Without medical indication	14.4	44.4
With medical indication	12.0	67.6
60 to 64 years	7.0	36.8
Without medical indication	4.8	29.5
With medical indication	2.1	53.0
65 years and older	19.4	61.3
Without medical indication	9.6	51.8
With medical indication	9.8	70.6
Total target population	34.6	52.6

* Persons under age 60 only belong to the target population if they have a medical indication.

Table 2 Size and influenza vaccine coverage by medical indication for vaccination (2019)

Medical indication	Size (% of target population)	Vaccine coverage (% of group)
Cardiovascular diseases	6.7	63.4
Pulmonary diseases	5.8	53.3
Diabetes mellitus	6.2	62.2
Weakened immune system	2.1	47.9
Chronic renal failure	1.7	63.3
Respiratory disorders caused by neurological conditions	1.7	56.8
HIV	0.1	44.9

Monitoring the Dutch National Influenza Prevention Program

An influenza infection is relatively harmless for most persons, however, in those who are vulnerable because of high age or medical conditions, influenza may cause serious complications. The World Health Organization (WHO) therefore advises that vulnerable groups are vaccinated. Vaccination reduces both the risk of getting an influenza infection and the risk of complications if someone is infected. The Dutch National Influenza Prevention Program was established in 1997 and coordinates and implements the influenza vaccination program. Persons at risk from complications due to (the consequences of) influenza are invited to visit their general practitioner for influenza vaccination free of charge. The target group is defined as all people aged 60 year or older and people with certain chronic diseases. The annual monitoring of vaccination uptake rates allows one to evaluate the number of people that are vaccinated (vaccine coverage) each year. In 2009, the European Union set a target to reach a vaccine coverage of 75% in the elderly and if possible in those with a medical indication.

About the monitor

The 2019 monitor was performed by NIVEL and commissioned by RIVM. Data from general practices participating in

NIVEL Primary Care Database were used for the analysis. For more information (in Dutch), see:
[Monitor Vaccinatiegraad Nationaal Programma Grieppreventie 2019. Utrecht: Nivel, 2020](#)

Citing this publication

The content may be used with acknowledgment of the source:

Heins M, Hooiveld M, Korevaar J. Vaccine Coverage Dutch National Influenza Prevention Program 2019: Brief monitor. Utrecht, Nivel: 2020.
